







## CBSE Class 12 Chemistry Question Paper Solution 2017



### Marking scheme – 2017



**CHEMISTRY (043)/ CLASS XII**



### Outside Delhi set (56/1)





Q No.	Value Points	Marks
1.	$\text{H}_3\text{PO}_4$	1
2.	2-Bromo-3-methylbut-2-en-1-ol	1
3.	a. Decreases b. No effect	$\frac{1}{2}$ $\frac{1}{2}$
4.		1
5.	Gel e.g. cheese, butter, jellies (any one)	$\frac{1}{2} + \frac{1}{2}$
6.	a. p-cresol < Phenol < p-nitrophenol b. 	1 1
	OR	
6	a.  b. 	1 1
7.	$n = \text{given mass} / \text{molar mass}$ $= 8.1 / 27 \text{ mol}$ $\text{Number of atoms} = \frac{8.1}{27} \times 6.022 \times 10^{23}$ $\text{Number of atoms in one unit cell} = 4 \text{ (fcc)}$ $\text{Number of unit cells} = \left[ \frac{8.1}{27} \times 6.022 \times 10^{23} \right] / 4$ $= 4.5 \times 10^{22}$ Or $27 \text{ g of Al contains} = 6.022 \times 10^{23} \text{ atoms}$ $8.1 \text{ g of Al contains} = (6.022 \times 10^{23} / 27) \times 8.1$ $\text{No of unit cells} = \text{total no of atoms} / 4$ $= \left[ \frac{8.1}{27} \times 6.022 \times 10^{23} \right] / 4$ $= 4.5 \times 10^{22}$	$\frac{1}{2}$ $\frac{1}{2}$  $\frac{1}{2}$ $\frac{1}{2}$  $\frac{1}{2}$ $\frac{1}{2}$  $\frac{1}{2}$ $\frac{1}{2}$


8.	<div><div></div><div></div></div> <p>a.)                      b.)</p>	1,1								
9.	Mercury cell Anode : $\text{Zn(Hg)} + 2\text{OH}^- \rightarrow \text{ZnO(s)} + \text{H}_2\text{O} + 2\text{e}^-$ Cathode : $\text{HgO} + \text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{Hg(l)} + 2\text{OH}^-$	1 $\frac{1}{2}$ $\frac{1}{2}$								
10.	(i) $\text{Na[Au(CN)}_2\text{]}$ (ii) $[\text{Pt(NH}_3)_4\text{Cl(NO}_2\text{)}]\text{SO}_4$	1 1								
11.	(a) Covalent solid / network solid , molecular solid (b) $\text{ZnO} \xrightarrow{\text{Heating}} \text{Zn}^{2+} + \frac{1}{2} \text{O}_2 + 2\text{e}^-$ Because excess $\text{Zn}^{2+}$ ions move to interstitial sites and the electrons move to neighbouring voids (c) Compounds prepared by combination of groups 12 and 16 behave like semiconductors. For eg ZnS, CdS , CdSe, HgTe (Any one)	$\frac{1}{2} + \frac{1}{2}$ 1 $\frac{1}{2} + \frac{1}{2}$								
12.	(a) $\Delta G^0 = -nFE_{\text{cell}}^0$ $n = 2$ $\Delta G^0 = -2 \times 96500 \text{ C/mol} \times 0.236 \text{ V}$ $= -45548 \text{ J/mol}$ $= -45.548 \text{ kJ/mol}$ (b) $Q = I t = 0.5 \times 2 \times 60 \times 60$ $= 3600 \text{ C}$ $96500 \text{ C} = 6.023 \times 10^{23} \text{ electrons}$ $3600 \text{ C} = 2.25 \times 10^{22} \text{ electrons}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1								
13.	(a) Linkage isomerism (b) In $[\text{NiCl}_4]^{2-}$ , due to the presence of $\text{Cl}^-$ , a weak field ligand no pairing occurs whereas in $[\text{Ni(CN)}_4]^{2-}$ , $\text{CN}^-$ is a strong field ligand and pairing takes place / diagrammatic representation (c) Because of very low CFSE which is not able to pair up the electrons.	1 1 1								
14.	<div><div>(a)</div><table><tr><td>Multimolecular colloid</td><td>Associated colloid</td></tr><tr><td>(a) Aggregation of large number of small atoms or molecules.</td><td>(a) Aggregation of large number of ions in concentrated solutions.</td></tr></table><div>(b)</div><table><tr><td>Coagulation</td><td>Peptization</td></tr><tr><td>(a) Settling down of colloidal particles.</td><td>(a) Conversion of precipitate into colloidal sol by adding small amount of</td></tr></table></div>	Multimolecular colloid	Associated colloid	(a) Aggregation of large number of small atoms or molecules.	(a) Aggregation of large number of ions in concentrated solutions.	Coagulation	Peptization	(a) Settling down of colloidal particles.	(a) Conversion of precipitate into colloidal sol by adding small amount of	1   1
Multimolecular colloid	Associated colloid									
(a) Aggregation of large number of small atoms or molecules.	(a) Aggregation of large number of ions in concentrated solutions.									
Coagulation	Peptization									
(a) Settling down of colloidal particles.	(a) Conversion of precipitate into colloidal sol by adding small amount of									

	electrolyte.	
	(c)	
	Homogenous catalysis	Heterogeneous catalysis
	(a) Reactants and catalyst are in same phase.	(a) Reactants and catalyst are in different phases.
		1
	OR	
14	(a) Dispersed phase-liquid , Dispersion medium – liquid (b) Both are surface phenomenon / both increase with increase in surface area (or any other correct similarity) (c) Hydrolysis / $\text{FeCl}_3 + 3\text{H}_2\text{O} \xrightarrow{\text{hydrolysis}} \text{Fe}(\text{OH})_3(\text{sol}) + 3\text{HCl}$	1 1 1
15.	$t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$ $20 \text{ min} = \frac{2.303}{k} \log \frac{100}{75} \quad \text{-(i)}$ $t = \frac{2.303}{k} \log \frac{100}{25} \quad \text{-(ii)}$ <p>Divide (i) equation by (ii)</p> $\frac{20}{t} = \frac{2.303}{k} \log \frac{100}{75} \div \frac{2.303}{k} \log \frac{100}{25}$ $= \frac{\log 4/3}{\log 4}$ $20/t = 0.1250/0.6021$ $t = 96.3 \text{ min}$ <p>(or any other correct procedure )</p>	$\frac{1}{2}$   $\frac{1}{2}$  $\frac{1}{2}$  $\frac{1}{2}$  1
16.	(i) 1- Bromopentane (ii) 2-Bromopentane (iii) 2-Bromo-2-methylbutane	1 1 1
17.	(a) Zone Refining – Impurities are more soluble in the melt than in the solid metal. (b) Mineral particles are wetted by oils forming froth while gangue particles are wetted by water and settle down. (c) Different components of a mixture are differently adsorbed on an adsorbent.	1 1 1
18.	(a) (A) $\text{CH}_3\text{CONH}_2$ (B) $\text{CH}_3\text{NH}_2$ (C) $\text{CH}_3\text{NC}$  (b) (A)   (B) 	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$  $\frac{1}{2}$  $\frac{1}{2}$

	(C)		½
19.	(a) $\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}_2$ , $\text{HOOC}-(\text{CH}_2)_4-\text{COOH}$ (b)  (c) $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$ , $\text{C}_6\text{H}_5-\text{CH}=\text{CH}_2$	and HCHO	1 1 1
20.	(a) Anionic detergents are sodium salts of sulphonated long chain alcohols or hydrocarbons / alkylbenzene sulphonate or detergents whose anionic part is involved in cleansing action. (b) Limited spectrum antibiotics are effective against a single organism or disease. (c) Antiseptics are the chemicals which either kill or prevent growth of microbes on living tissues.		1 1 1
21.	(a) Red phosphorous being polymeric is less reactive than white phosphorous which has discrete tetrahedral structure. (b) They readily accept an electron to attain noble gas configuration. (c) Because of higher oxidation state(+5) of nitrogen in $\text{N}_2\text{O}_5$		1 1 1
22.	(i) Due to the resonance, the electron pair of nitrogen atom gets delocalised towards carbonyl group / resonating structures. (ii) Because of +I effect in methylamine electron density at nitrogen increases whereas in aniline resonance takes place and electron density on nitrogen decreases / resonating structures. (iii) Due to protonation of aniline / formation of anilinium ion		1 1 1
23.	(i) Concerned, caring, socially alert, leadership (or any other 2 values) (ii) Starch (iii) $\alpha$ -Helix and $\beta$ -pleated sheets (iv) Vitamin B / $\text{B}_1$ / $\text{B}_2$ / $\text{B}_6$ / C (any two)		½ + ½ 1 ½ + ½ ½ + ½
24.	a. (i) Availability of partially filled d-orbitals / comparable energies of ns and (n-1) d orbitals (ii) Completely filled d-orbitals / absence of unpaired d electrons cause weak metallic bonding (iii) Because $\text{Mn}^{2+}$ has $d^5$ as a stable configuration whereas $\text{Cr}^{3+}$ is more stable due to stable $t_{2g}^3$ b) Similarity-both are stable in +3 oxidation state/ both show contraction/ irregular electronic configuration (or any other suitable similarity) Difference- actinoids are radioactive and lanthanoids are not / actinoids show wide range of oxidation states but lanthanoids don't (or any other correct difference)		1 1 1 1 1
	OR		
24	a. (i) $\text{Cr}^{3+}$ , half filled $t_{2g}^3$ (ii) $\text{Mn}^{3+}$ , due to stable $d^5$ configuration in $\text{Mn}^{2+}$		½ + ½ ½ + ½




	(iii) $Ti^{4+}$ , No unpaired electrons b. (i) $2MnO_4^- + 16H^+ + 5S^{2-} \rightarrow 5S + 2Mn^{2+} + 8H_2O$ (ii) $2KMnO_4 \rightarrow K_2MnO_4 + MnO_2 + O_2$	$\frac{1}{2} + \frac{1}{2}$ 1 1				
25	a) $\Delta T_f = K_f m$ Here, $m = w_2 \times 1000 / M_2 \times M_1$ $273.15 - 269.15 = K_f \times 10 \times 1000 / 342 \times 90$ $K_f = 12.3 \text{ K kg/mol}$ $\Delta T_f = K_f m$ $= 12.3 \times 10 \times 1000 / 180 \times 90$ $= 7.6 \text{ K}$ $T_f = 273.15 - 7.6 = 265.55 \text{ K}$ (or any other correct method) b) (i) Number of moles of solute dissolved in per kilo gram of the solvent. (ii) Abnormal molar mass: If the molar mass calculated by using any of the colligative properties to be different than theoretically expected molar mass.	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1 1				
	OR					
25.	(a) $(P_A^0 - P_A) / P_A^0 = (w_B \times M_A) / (M_B \times w_A)$ $\frac{23.8 - P_A}{23.8} = (30 \times 18) / 60 \times 846$ $23.8 - P_A = 23.8 \times [(30 \times 18) / 60 \times 846]$ $23.8 - P_A = 0.2532$ $P_A = 23.55 \text{ mm Hg}$ (b)	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1				
	<table><tr><th>Ideal solution</th><th>Non ideal solution</th></tr><tr><td>(a) It obeys Raoult's law over the entire range of concentration. (b) <math>\Delta_{mix} H = 0</math> (c) <math>\Delta_{mix} V = 0</math></td><td>(a) Does not obey Raoult's law over the entire range of concentration. (b) <math>\Delta_{mix} H</math> is not equal to 0. (c) <math>\Delta_{mix} V</math> is not equal to 0.</td></tr></table> (any two correct difference)	Ideal solution	Non ideal solution	(a) It obeys Raoult's law over the entire range of concentration. (b) $\Delta_{mix} H = 0$ (c) $\Delta_{mix} V = 0$	(a) Does not obey Raoult's law over the entire range of concentration. (b) $\Delta_{mix} H$ is not equal to 0. (c) $\Delta_{mix} V$ is not equal to 0.	1 + 1
Ideal solution	Non ideal solution					
(a) It obeys Raoult's law over the entire range of concentration. (b) $\Delta_{mix} H = 0$ (c) $\Delta_{mix} V = 0$	(a) Does not obey Raoult's law over the entire range of concentration. (b) $\Delta_{mix} H$ is not equal to 0. (c) $\Delta_{mix} V$ is not equal to 0.					
26.	a.  (i)  (ii)	1 1				

	<p>(iii) <math>\text{CH}_3\text{-CH=CH-CHO}</math></p> <p>b. (i) Tollen's reagent test: Add ammoniacal solution of silver nitrate (Tollen's Reagent) in both the solutions. Butanal gives silver mirror whereas Butan-2-one does not.</p> <p>(ii) Add neutral <math>\text{FeCl}_3</math> in both the solutions, phenol forms violet colour but benzoic acid does not.</p> <p>(or any other correct test)</p>	<p>1</p> <p>1</p> <p>1</p>
	OR	
26	<p>(a) (i) Étard reaction</p> <p></p> <p></p> <p>(ii) Stephen reaction</p> <p><math>\text{RCN} + \text{SnCl}_2 + \text{HCl} \longrightarrow \text{RCH=NH} \xrightarrow{\text{H}_3\text{O}^+} \text{RCHO}</math></p> <p>Or</p> <p><math>\text{RCN} \xrightarrow[\text{(ii) H}_3\text{O}^+]{\text{(i) SnCl}_2 + \text{HCl}} \text{RCHO}</math></p> <p>(b) (i)</p> <p></p> <p>(ii)</p> <p></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>




1	Dr. (Mrs.) Sangeeta Bhatia		12	Sh. S. Vallabhan	
2	Dr. K.N. Uppadhya		13	Dr. Bhagyabati Nayak	
3	Prof. R.D. Shukla		14	Ms. Anila Mechur Jayachandran	
4	Sh. S.K. Munjal		15	Mrs. Deepika Arora	
5	Sh. D.A. Mishra		16	Ms. Seema Bhatnagar	
6	Sh. Rakesh Dhawan		17	Mrs. Sushma Sachdeva	
7	Dr. (Mrs.) Sunita Ramrakhiani		18	Dr. Azhar Aslam Khan	
8	Mrs. Preeti Kiran		19	Mr. Roop Narain Chauhan	
9	Ms. Neeru Sofat		20	Mr. Mukesh Kumar Kaushik	
10	Sh. Pawan Singh Meena		21	Ms. Abha Chaudhary	
11	Mrs. P. Nirupama Shankar		22	Ms. Garima Bhutani	

**Marking scheme – 2017**  
**CHEMISTRY (043)/ CLASS XII**  
**Outside Delhi set (56/2)**

Q.No	Value points	Marks
1.	a. Decreases b. No change	$\frac{1}{2}$ $\frac{1}{2}$
2.	Sol : example- paints, cell fluids (any one)	$\frac{1}{2} + \frac{1}{2}$
3.	3-phenyl-prop-2-en-1-ol	1
4.	H <sub>2</sub> SO <sub>4</sub>	1
5.		1
6.	(i) [Cr(en) <sub>3</sub> ]Cl <sub>3</sub> (ii) K <sub>2</sub> [Zn(OH) <sub>4</sub> ]	1 1
7.	(a)   (b) 	1  1
8.	Lead storage battery Anode : Pb <sub>(s)</sub> + SO <sub>4</sub> <sup>2-</sup> <sub>(aq)</sub> → PbSO <sub>4(s)</sub> + 2e <sup>-</sup> Cathode : PbO <sub>2</sub> + SO <sub>4</sub> <sup>2-</sup> <sub>(aq)</sub> + 4H <sup>+</sup> + 2e <sup>-</sup> → PbSO <sub>4(s)</sub> + 2 H <sub>2</sub> O <sub>(l)</sub>	1 $\frac{1}{2}$ $\frac{1}{2}$
9.	n = given mass / molar mass = 8.1 / 27 mol Number of atoms = $\frac{8.1}{27} \times 6.022 \times 10^{23}$ Number of atoms in one unit cell = 4 (fcc) Number of unit cells = $\left[ \frac{8.1}{27} \times 6.022 \times 10^{23} \right] / 4$ = 4.5 × 10 <sup>22</sup>	$\frac{1}{2}$ $\frac{1}{2}$   $\frac{1}{2}$ $\frac{1}{2}$



13.	<p>(a) <math>\Delta G^0 = -nFE^0_{\text{cell}}</math>  <math>n = 2</math>  <math>\Delta G^0 = -2 \times 96500 \text{ C/mol} \times 0.236 \text{ V}</math>  <math>= -45548 \text{ J/mol}</math>  <math>= -45.548 \text{ kJ/mol}</math></p> <p>(b) <math>Q = I t = 0.5 \times 2 \times 60 \times 60</math>  <math>= 3600 \text{ C}</math>  <math>96500 \text{ C} = 6.023 \times 10^{23} \text{ electrons}</math>  <math>3600 \text{ C} = 2.25 \times 10^{22} \text{ electrons}</math></p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p>1</p>
14.	<p>(i) Due to the resonance, the electron pair of nitrogen atom gets delocalised towards carbonyl group / resonating structures.</p> <p>(ii) Because of +I effect in methylamine electron density at nitrogen increases whereas in aniline resonance takes place and electron density on nitrogen decreases / resonating structures.</p> <p>(iii) Due to protonation of aniline / formation of anilinium ion</p>	<p>1</p> <p>1</p> <p>1</p>
15	<p>(a) Red phosphorous being polymeric is less reactive than white phosphorous which has discrete tetrahedral structure.</p> <p>(b) They readily accept an electron to attain noble gas configuration.</p> <p>(c) Because of higher oxidation state(+5) of nitrogen in <math>\text{N}_2\text{O}_5</math></p>	<p>1</p> <p>1</p> <p>1</p>
16	<p>(a) Anionic detergents are sodium salts of sulphonated long chain alcohols or hydrocarbons / alkylbenzene sulphonate or detergents whose anionic part is involved in cleansing action.</p> <p>(b) Narrow spectrum antibiotics- which are effective against either gram positive or gram negative bacteria.</p> <p>(c) Chemical compounds which are used for the treatment of excess acid produced in the stomach.</p>	<p>1</p> <p>1</p> <p>1</p>
17	<p>(a) <math>\text{CH}_2=\text{CHCl}</math></p> <p>(b)  and <math>\text{HCHO}</math></p> <p>(c) <math>\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2</math>, <math>\text{CH}_2=\text{CHCN}</math></p>	<p>1</p> <p>1</p> <p>1</p>
18.	<p>(i) 1-Bromopentane</p> <p>(ii) 2-Bromopentane</p> <p>(iii) 2-Bromo-2-methylbutane</p>	<p>1</p> <p>1</p> <p>1</p>

19.	$t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$ $20 \text{ min} = \frac{2.303}{k} \log \frac{100}{75} \quad \text{--- (i)}$ $t = \frac{2.303}{k} \log \frac{100}{25} \quad \text{--- (ii)}$ <p>Divide (i) equation by (ii)</p> $\frac{20}{t} = \frac{2.303}{k} \log \frac{100}{75}$ <hr/> $\frac{2.303}{k} \log \frac{100}{25}$ $= \frac{\log 4/3}{\log 4}$ $20/t = 0.1250/0.6021$ $t = 96.3 \text{ min}$ <p>(or any other correct procedure )</p>	$\frac{1}{2}$  $\frac{1}{2}$  $\frac{1}{2}$  $\frac{1}{2}$  1												
20	<p>(a)</p> <table> <tr> <td>Multimolecular colloid</td> <td>Associated colloid</td> </tr> <tr> <td>(a) Aggregation of large number of small atoms or molecules.</td> <td>(a) Aggregation of large number of ions in concentrated solutions.</td> </tr> </table> <p>(b)</p> <table> <tr> <td>Coagulation</td> <td>Peptization</td> </tr> <tr> <td>(a) Settling down of colloidal particles.</td> <td>(a) Conversion of precipitate into colloidal sol by adding small amount of electrolyte.</td> </tr> </table> <p>(c)</p> <table> <tr> <td>Homogenous catalysis</td> <td>Heterogeneous catalysis</td> </tr> <tr> <td>(a) Reactants and catalyst are in same phase.</td> <td>(a) Reactants and catalyst are in different phases.</td> </tr> </table>	Multimolecular colloid	Associated colloid	(a) Aggregation of large number of small atoms or molecules.	(a) Aggregation of large number of ions in concentrated solutions.	Coagulation	Peptization	(a) Settling down of colloidal particles.	(a) Conversion of precipitate into colloidal sol by adding small amount of electrolyte.	Homogenous catalysis	Heterogeneous catalysis	(a) Reactants and catalyst are in same phase.	(a) Reactants and catalyst are in different phases.	1   <
Multimolecular colloid	Associated colloid													
(a) Aggregation of large number of small atoms or molecules.	(a) Aggregation of large number of ions in concentrated solutions.													
Coagulation	Peptization													
(a) Settling down of colloidal particles.	(a) Conversion of precipitate into colloidal sol by adding small amount of electrolyte.													
Homogenous catalysis	Heterogeneous catalysis													
(a) Reactants and catalyst are in same phase.	(a) Reactants and catalyst are in different phases.													