EXERCISE 8.1

PAGE NO: 166

Expand each of the expressions in Exercises 1 to 5.

1. $(1 - 2x)^5$

Solution:

From binomial theorem expansion we can write as

$$(1-2x)^{5} = {}^{5}C_{0}(1)^{5} - {}^{5}C_{1}(1)^{4}(2x) + {}^{5}C_{2}(1)^{3}(2x)^{2} - {}^{5}C_{3}(1)^{2}(2x)^{3} + {}^{5}C_{4}(1)^{1}(2x)^{4} - {}^{5}C_{5}(2x)^{5} = 1 - 5(2x) + 10(4x)^{2} - 10(8x^{3}) + 5(16x^{4}) - (32x^{5}) = 1 - 10x + 40x^{2} - 80x^{3} - 32x^{5}$$

2. $\left(\frac{2}{x} - \frac{x}{2}\right)^5$

Solution:

From binomial theorem, given equation can be expanded as

$$\begin{pmatrix} \frac{2}{x} - \frac{x}{2} \end{pmatrix}^5 = {}^5 \operatorname{C}_0 \left(\frac{2}{x}\right)^3 - {}^5 \operatorname{C}_1 \left(\frac{2}{x}\right)^4 \left(\frac{x}{2}\right) + {}^5 \operatorname{C}_2 \left(\frac{2}{x}\right)^3 \left(\frac{x}{2}\right)^2 \\ - {}^3 \operatorname{C}_3 \left(\frac{2}{x}\right)^2 \left(\frac{x}{2}\right)^3 + {}^3 \operatorname{C}_4 \left(\frac{2}{x}\right) \left(\frac{x}{2}\right)^4 - {}^3 \operatorname{C}_5 \left(\frac{x}{2}\right)^5 \\ = \frac{32}{x^5} - 5 \left(\frac{16}{x^4}\right) \left(\frac{x}{2}\right) + 10 \left(\frac{8}{x^3}\right) \left(\frac{x^2}{4}\right) - 10 \left(\frac{4}{x^2}\right) + 5 \left(\frac{2}{x}\right) \left(\frac{x^4}{16}\right) - \frac{x^5}{32} \\ = \frac{32}{x^5} - \frac{40}{x^3} + \frac{20}{x} - 5x + \frac{5}{8}x^3 - \frac{x^3}{32}$$

3. (2x – 3)⁶

Solution:

From binomial theorem, given equation can be expanded as

$$egin{aligned} &(2x-3)^6 = ^6\mathrm{C}_0(2x)^6 - ^6\mathrm{C}_1(2x)^5(3) + ^6\mathrm{C}_1(2x)^4(3)^2 - ^4\mathrm{C}_3(2x)^3(3)^3\ &= 64x^6 - 6\left(32x^5
ight)(3) + 15\left(16x^4
ight)(9) - 20\left(8x^3
ight)(27)\ &+ 15\left(4x^2
ight)(81) - 6(2x)(243) + 729\ &= 64x^6 - 576x^5 + 2160x^4 - 4320x^3 + 4860x^2 - 2916x + 729 \end{aligned}$$

$$4. \left(\frac{x}{3} + \frac{1}{x}\right)^5$$

Solution:

From binomial theorem, given equation can be expanded as

$$\left(\frac{x}{3} + \frac{1}{x}\right)^5 = {}^5 C_0 \left(\frac{x}{3}\right)^5 + {}^3 C_1 \left(\frac{x}{3}\right)^4 \left(\frac{1}{x}\right) + {}^3 C_2 \left(\frac{x}{3}\right)^3 \left(\frac{1}{x}\right)^2$$

$$= \frac{x^5}{243} + 5 \left(\frac{x^4}{81}\right) \left(\frac{1}{x}\right) + 10 \left(\frac{x^3}{27}\right) \left(\frac{1}{x^2}\right) + 10 \left(\frac{x^2}{9}\right) \left(\frac{1}{x^3}\right) + 5 \left(\frac{x}{3}\right) \left(\frac{1}{x^4}\right) + \frac{1}{x^5}$$

$$= \frac{x^5}{243} + \frac{5x^3}{81} + \frac{10x}{27} + \frac{10}{9x} + \frac{5}{3x^3} + \frac{1}{x^3}$$

5.
$$\left(x+\frac{1}{x}\right)^6$$

Solution:

From binomial theorem, given equation can be expanded as

$$\begin{pmatrix} \mathbf{x} + \frac{1}{\mathbf{x}} \end{pmatrix}^6 = {}^6 \mathbf{C}_0(\mathbf{x})^6 + {}^6 \mathbf{C}_1(\mathbf{x})' \left(\frac{1}{\mathbf{x}}\right) + {}^6 \mathbf{C}_2(\mathbf{x})^4 \left(\frac{1}{\mathbf{x}}\right)^2 + {}^6 \mathbf{C}_3(\mathbf{x})^3 \left(\frac{1}{\mathbf{x}}\right)^3 + {}^6 \mathbf{C}_4(\mathbf{x})^2 \left(\frac{1}{\mathbf{x}}\right)^4 + {}^6 \mathbf{C}_3(\mathbf{x}) \left(\frac{1}{\mathbf{x}}\right)^5 + {}^6 \mathbf{C}_6 \left(\frac{1}{\mathbf{x}}\right)^6 = \mathbf{x}^4 + 6(\mathbf{x})^3 \left(\frac{1}{\mathbf{x}}\right) + 15(\mathbf{x})^4 \left(\frac{1}{\mathbf{x}^2}\right) + 20(\mathbf{x})^3 \left(\frac{1}{\mathbf{x}^3}\right) + 15(\mathbf{x})^2 \left(\frac{1}{\mathbf{x}^4}\right) + 6(\mathbf{x}) \left(\frac{1}{\mathbf{x}^5}\right) + \frac{1}{\mathbf{x}^6} \\ = \mathbf{x}^6 + 6\mathbf{x}^4 + 15\mathbf{x}^2 + 20 + \frac{15}{\mathbf{x}^2} + \frac{6}{\mathbf{x}^4} + \frac{1}{\mathbf{x}^6}$$

6. (96)³

Solution:

NCERT Solutions for Class 11Maths Chapter 8 **Binomial Theorem**

Given (96)³

96 can be expressed as the sum or difference of two numbers and then binomial theorem can be applied.

The given question can be written as 96 = 100 - 4

 $(96)^3 = (100 - 4)^3$

```
= {}^{3}C_{0} (100)^{3} - {}^{3}C_{1} (100)^{2} (4) - {}^{3}C_{2} (100) (4)^{2} - {}^{3}C_{3} (4)^{3}
```

```
= (100)^3 - 3 (100)^2 (4) + 3 (100) (4)^2 - (4)^3
```

```
= 1000000 - 120000 + 4800 - 64
```

```
= 884736
```

7. (102)⁵

Solution:

Given (102)⁵

102 can be expressed as the sum or difference of two numbers and then binomial theorem can be applied.

The given question can be written as 102 = 100 + 2

 $(102)^5 = (100 + 2)^5$

$$= {}^{5}C_{0} (100)^{5} + {}^{5}C_{1} (100)^{4} (2) + {}^{5}C_{2} (100)^{3} (2)^{2} + {}^{5}C_{3} (100)^{2} (2)^{3} + {}^{5}C_{4} (100) (2)^{4} + {}^{5}C_{5} (2)^{5}$$

 $= (100)^{5} + 5 (100)^{4} (2) + 10 (100)^{3} (2)^{2} + 5 (100) (2)^{3} + 5 (100) (2)^{4} + (2)^{5}$

= 100000000 + 100000000 + 40000000 + 80000 + 8000 + 32

= 11040808032

8. (101)⁴

Solution:

Given (101)⁴

101 can be expressed as the sum or difference of two numbers and then binomial theorem can be applied.

The given question can be written as 101 = 100 + 1 $(101)^4 = (100 + 1)^4$ $= {}^{4}C_{0}(100)^{4} + {}^{4}C_{1}(100)^{3}(1) + {}^{4}C_{2}(100)^{2}(1)^{2} + {}^{4}C_{3}(100)(1)^{2} + {}^{4}C_{4}(1)^{4}$ $= (100)^4 + 4 (100)^3 + 6 (100)^2 + 4 (100) + (1)^4$

= 10000000 + 400000 + 60000 + 400 + 1



= 1040604001

```
9. (99)<sup>5</sup>
```

Solution:

Given (99)⁵

99 can be written as the sum or difference of two numbers then binomial theorem can be applied.

The given question can be written as 99 = 100 -1

 $(99)^5 = (100 - 1)^5$

$$= {}^{5}C_{0} (100)^{5} - {}^{5}C_{1} (100)^{4} (1) + {}^{5}C_{2} (100)^{3} (1)^{2} - {}^{5}C_{3} (100)^{2} (1)^{3} + {}^{5}C_{4} (100) (1)^{4} - {}^{5}C_{5} (1)^{5}$$

 $= (100)^{5} - 5 (100)^{4} + 10 (100)^{3} - 10 (100)^{2} + 5 (100) - 1$

- = 100000000 500000000 + 10000000 100000 + 500 1
- = 9509900499

10. Using Binomial Theorem, indicate which number is larger (1.1)¹⁰⁰⁰⁰ or 1000.

Solution:

By splitting the given 1.1 and then applying binomial theorem, the first few terms of

 $(1.1)^{10000}$ can be obtained as

 $(1.1)_{10000} = (1 + 0.1)_{10000}$

 $= (1 + 0.1)^{10000} C_1(1.1) + other positive terms$

= $1 + 10000 \times 1.1 + other positive terms$

- = 1 + 11000 + other positive terms
- > 1000
- $(1.1)^{10000} > 1000$

11. Find $(a + b)^4 - (a - b)^4$. Hence, evaluate $(\sqrt{3} + \sqrt{2})^4 - (\sqrt{3} - \sqrt{2})^4$.

Solution:

Using binomial theorem the expression $(a + b)^4$ and $(a - b)^4$, can be expanded $(a + b)^4 = {}^4C_0 a^4 + {}^4C_1 a^3 b + {}^4C_2 a^2 b^2 + {}^4C_3 a b^3 + {}^4C_4 b^4$

$$(a - b)^{4} = {}^{4}C_{0} a^{4} - {}^{4}C_{1} a^{3} b + {}^{4}C_{2} a^{2} b^{2} - {}^{4}C_{3} a b^{3} + {}^{4}C_{4} b^{4}$$
Now $(a + b)^{4} - (a - b)^{4} = {}^{4}C_{0} a^{4} + {}^{4}C_{1} a^{3} b + {}^{4}C_{2} a^{2} b^{2} + {}^{4}C_{3} a b^{3} + {}^{4}C_{4} b^{4} - [{}^{4}C_{0} a^{4} - {}^{4}C_{1} a^{3} b + {}^{4}C_{2} a^{2} b^{2} - {}^{4}C_{3} a b^{3} + {}^{4}C_{4} b^{4}]$

$$= 2 ({}^{4}C_{1} a^{3} b + {}^{4}C_{3} a b^{3})$$

$$= 2 ({}^{4}C_{1} a^{3} b + {}^{4}C_{3} a b^{3})$$

$$= 2 ({}^{4}a^{3} b + {}^{4}ab^{3})$$

$$= 8ab (a^{2} + b^{2})$$
Now by substituting a = $\sqrt{3}$ and b = $\sqrt{2}$ we get
 $(\sqrt{3} + \sqrt{2})^{4} - (\sqrt{3} - \sqrt{2})^{4} = 8 (\sqrt{3}) (\sqrt{2}) \{(\sqrt{3})^{2} + (\sqrt{2})^{2}\}$

$$= 8 (\sqrt{6}) (3 + 2)$$

$$= 40 \sqrt{6}$$

12. Find $(x + 1)^6 + (x - 1)^6$. Hence or otherwise evaluate $(\sqrt{2} + 1)^6 + (\sqrt{2} - 1)^6$

Solution:

Using binomial theorem the expressions, $(x + 1)^6$ and $(x - 1)^6$ can be expressed as $(x + 1)^6 = {}^6C_0 x^6 + {}^6C_1 x^5 + {}^6C_2 x^4 + {}^6C_3 x^3 + {}^6C_4 x^2 + {}^6C_5 x + {}^6C_6$ $(x - 1)^6 = {}^6C_0 x^6 - {}^6C_1 x^5 + {}^6C_2 x^4 - {}^6C_3 x^3 + {}^6C_4 x^2 - {}^6C_5 x + {}^6C_6$ Now, $(x + 1)^6 - (x - 1)^6 = {}^6C_0 x^6 + {}^6C_1 x^5 + {}^6C_2 x^4 + {}^6C_3 x^3 + {}^6C_4 x^2 + {}^6C_5 x + {}^6C_6 - [{}^6C_0 x^6 - {}^6C_1 x^5 + {}^6C_2 x^4 - {}^6C_3 x^3 + {}^6C_4 x^2 - {}^6C_5 x + {}^6C_6]$ $= 2 [{}^6C_0 x^6 + {}^6C_2 x^4 + {}^6C_4 x^2 + {}^6C_6]$ $= 2 [x^6 + 15x^4 + 15x^2 + 1]$ Now by substituting x = V2 we get $(V2 + 1)^6 - (V2 - 1)^6 = 2 [(V2)^6 + 15(V2)^4 + 15(V2)^2 + 1]$ $= 2 (8 + 15 \times 4 + 15 \times 2 + 1)$ = 2 (8 + 60 + 30 + 1) = 2 (99)= 198

13. Show that $9^{n+1} - 8n - 9$ is divisible by 64, whenever n is a positive integer.

Solution:

In order to show that $9^{n+1} - 8n - 9$ is divisible by 64, it has to be show that $9^{n+1} - 8n - 9 =$

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

64 k, where k is some natural number Using binomial theorem, $(1 + a)^m = {}^mC_0 + {}^mC_1 a + {}^mC_2 a^2 + + {}^mC_m a^m$ For a = 8 and m = n + 1 we get $(1 + 8)_{n+1} = {}_{n+1}C_0 + {}_{n+1}C_1 (8) + {}_{n+1}C_2 (8)_2 + + {}_{n+1}C_{n+1} (8)_{n+1}$ $9^{n+1} = 1 + (n + 1) 8 + 8^2 [{}^{n+1}C_2 + {}^{n+1}C_3 (8) + + {}^{n+1}C_{n+1} (8)_{n-1}]$ $9_{n+1} = 9 + 8n + 64 [{}_{n+1}C_2 + {}_{n+1}C_3 (8) + + {}^{n+1}C_{n+1} (8)_{n-1}]$ $9^{n+1} - 8n - 9 = 64 k$ Where k = $[{}^{n+1}C_2 + {}^{n+1}C_3 (8) + + {}^{n+1}C_{n+1} (8)_{n-1}]$ is a natural number. Thus, $9^{n+1} - 8n - 9$ is divisible by 64, whenever n is positive integer. Hence the proof

14. Prove that

 $\sum_{r=0}^{n} 3^{r \ n} C_r = 4^n$

Solution:

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

By Binomial Theorem

$$\sum_{r=0}^{n} {n \choose r} a^{n-r} b^r = (a+b)^n$$

On right side we need 4^n so we will put the values as, Putting b = 3 & a = 1 in the above equation, we get

$$\sum_{r=0}^{n} {n \choose r} (1)^{n-r} (3)^{r} = (1+3)^{n}$$
$$\sum_{r=0}^{n} {n \choose r} (1)(3)^{r} = (4)^{n}$$
$$\sum_{r=0}^{n} {n \choose r} (3)^{r} = (4)^{n}$$

Hence Proved.



NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

EXERCISE 8.2

PAGE NO: 171

Find the coefficient of 1. x^5 in $(x + 3)^8$

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here x^{5} is the T_{r+1} term so a = x, b = 3 and n = 8 $T_{r+1} = {}^{8}C_{r} x^{8-r} 3^{r}$(i) For finding out x^{5} We have to equate $x^{5} = x^{8-r} \Rightarrow$ r = 3Putting value of r in (i) we get $T_{3+1} = {}^{8}C_{3} x^{8-3} 3^{3}$ $T_{4} = \frac{8!}{3! 5!} \times x^{5} \times 27$

= 1512 x⁵

Hence the coefficient of $x^5 = 1512$

2. $a^{5}b^{7}$ in $(a - 2b)^{12}$.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here a = a, b = -2b & n = 12Substituting the values, we get $T_{r+1} = {}^{12}C_{r} a^{12-r} (-2b)^{r}$(i) To find a^{5} We equate $a^{12-r} = a^{5} r$ = 7Putting r = 7 in (i) $T_{8} = {}^{12}C_{7} a^{5} (-2b)^{7}$



$$T_8 = \frac{12!}{7!5!} \times a^5 \times (-2)^7 b^7$$

= -101376 a⁵ b⁷
Hence the coefficient of a⁵b⁷= -101376

Write the general term in the expansion of 3. $(x^2 - y)^6$

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = n C r a_{n-r} b_{r}$(i) Here $a = x^2$, n = 6 and b = -yPutting values in (i) $T_{r+1} = 6Cr \times 2(6-r) (-1)r yr$ $= \frac{6!}{r! (6-r)!} \times x^{12-2r} \times (-1)^r \times y^r$ $= -1^r \frac{6!}{r! (6-r)!} \times x^{12-2r} \times y^r$ $= -1_r 6Cr . X_{12-2r} . yr$

4. $(x^2 - y x)^{12}$, $x \neq 0$.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here n = 12, a= x² and b = -y x Substituting the values we get $T_{n+1} = {}_{12}C_{r} \times x_{2(12-r)} (-1)_{r} y_{r} x_{r}$ $= \frac{12!}{r!(12-r)!} \times x^{24-2r} - 1^{r} y^{r} x^{r}$

$$= \frac{-1^{r}}{r!(12-r)!} x^{24-r} y^{r}$$

 $= -1_{r_{12}C_r} \cdot X_{24} - 2r \cdot y_r$

5. Find the 4th term in the expansion of $(x - 2y)^{12}$.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$

Here a = x, n = 12, r = 3 and b = -2y

By substituting the values we get

$$T_{4} = {}^{12}C_{3} x^{9} (-2y)^{3}$$

= $\frac{12!}{3!9!} \times x^{9} \times -8 \times y^{3}$
= $-\frac{12 \times 11 \times 10 \times 8}{3 \times 2 \times 1} \times x^{9} y^{3}$
= $-1760 x^{9} y^{3}$

6. Find the 13th term in the expansion of

$$\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}, x \neq 0$$

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$

Here a=9x,
$$b = -\frac{1}{3\sqrt{x}} = -\frac{1}{3\sqrt{x}}$$
 n =18 and r = 12

Putting values

$$T_{13} = \frac{18!}{12! \, 6!} \, 9x^{18-12} \left(-\frac{1}{3\sqrt{x}} \right)^{12}$$
$$= \frac{(18 \times 17 \times 16 \times 15 \times 14 \times 13 \times 12!)}{12! \times 6 \times 5 \times 4 \times 3 \times 2 \times 1} \times 3^{12} \times x^6 \times \frac{1}{x^6} \times \frac{1}{3^{12}}$$
$$= 18564$$

Find the middle terms in the expansions of

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

7.
$$\left(3-\frac{x^3}{6}\right)^7$$

Solution:

Here n = 7 so there would be two middle terms given by

 $\left(\frac{n+1}{2}^{th}\right)$ term = 4 th and $\left(\frac{n+1}{2}+1\right)$ th term = 5th

We have

a = 3, n = 7 and b =
$$-\frac{x^3}{6}$$

For T₄, r= 3

The term will be

$$T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$$

$$T_{4} = \frac{7!}{3!} 3^{4} \left(-\frac{x^{3}}{6}\right)^{3}$$
$$= -\frac{7 \times 6 \times 5 \times 4}{3 \times 2 \times 1} \times 3^{4} \times \frac{x^{9}}{2^{3} 3^{3}}$$
$$= -\frac{105}{8} x^{9}$$

For T_5 term, r = 4

The term T_{r+1} in the binomial expansion is given by

$$T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$$

$$T_{5} = \frac{7!}{4! \, 3!} \, 3^{3} \left(-\frac{x^{3}}{6}\right)^{4}$$

$$= \frac{7 \times 6 \times 5 \times 4!}{4! \, 3!} \times \frac{3^{3}}{2^{4} 3^{4}} \times x^{3} = \frac{35 \, x^{12}}{48}$$

8.
$$\left(\frac{x}{3}+9y\right)^{10}$$

Solution:

Here n is even so the middle term will be given by $\left(\frac{n+1}{2}\right)^{\text{th}}$ term = 6th term

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$

Now
$$a = \frac{x}{3}$$
, $b = 9y, n = 10$ and $r = 5$

Substituting the values

$$T_{6} = \frac{10!}{5! \, 5!} \times \left(\frac{x}{3}\right)^{5} \times (9y)^{5}$$
$$= \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2 \times 1} \times \frac{x^{5}}{3^{5}} \times 3^{10} \times y^{5}$$
$$= 61236 \, x^{5} y^{5}$$

9. In the expansion of $(1 + a)^{m+n}$, prove that coefficients of a^m and a^n are equal.

Solution:

We know that the general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here n= m+n, a = 1 and b= a Substituting the values in the general form $T_{r+1} = {}^{m+n}C_{r} 1{}^{m+n-r}a^{r}$

$$= {}^{m+n} C_r a^r$$
.....(i)

Now we have that the general term for the expression is,

 $T_{r+1} = m+n Cr ar$ Now, For coefficient of a^m $T_{m+1} = m+n Cm am$ Hence, for coefficient of a^m , value of r = m

So, the coefficient is ^{m+n} C_m Similarly, Coefficient of aⁿ is ^{m+n} C_n ${}^{m+n}C_{m} = \frac{(m+n)!}{m!n!}$ And also, ^{m+n} C_n = $\frac{(m+n)!}{m!n!}$

(m+n)!

The coefficient of a^m and aⁿ are same that is mini

10. The coefficients of the $(r - 1)^{th}$, r^{th} and $(r + 1)^{th}$ terms in the expansion of $(x + 1)^n$ are in the ratio 1 : 3 : 5. Find n and r.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here the binomial is $(1+x)^{n}$ with a = 1, b = x and n = nThe $(r+1)^{th}$ term is given by $T_{(r+1)} = {}^{n}C_{r} 1_{n-r} x_{r}$ $T_{(r+1)} = {}^{n}C_{r} x_{r}$ The coefficient of $(r+1)^{th}$ term is ${}^{n}C_{r}$ The rth term is given by $(r-1)^{th}$ term $T_{(r+1-1)} = {}^{n}C_{r-1} x_{r-1}$ $T_{r} = {}^{n}C_{r-1} x_{r-1}$... the coefficient of rth term is ${}^{n}C_{r-1}$ For $(r-1)^{th}$ term we will take $(r-2)^{th}$ term $T_{r-2+1} = {}^{n}C_{r-2} x_{r-2}$ $T_{r-1} = {}^{n}C_{r-2} x_{r-2}$... the coefficient of $(r-1)^{th}$, rth and r+1th term are in ratio 1:3:5 Therefore,



 $\frac{\text{the coefficient of } r - 1^{\text{th}} \text{ term}}{\text{coefficient of } r^{\text{th}} \text{ term}} = \frac{1}{3}$ $n_{\substack{\substack{c \\ \frac{r-2}{n c \\ r-1}}}} = \frac{1}{3}$ $\Rightarrow \frac{\frac{n!}{(r-2)!(n-r+2)!}}{\frac{n!}{(r-1)!(n-r+1)!}} = \frac{1}{3}$

On rearranging we get

$$\frac{n!}{(r-2)!(n-r+2)!} \times \frac{(r-1)!(n-r+1)!}{n!} = \frac{1}{3}$$

By multiplying

$$\Rightarrow \frac{(r-1)(r-2)!(n-r+1)!}{(r-2)!(n-r+2)!} = \frac{1}{3}$$
$$\Rightarrow \frac{(r-1)(n-r+1)!}{(n-r+2)(n-r+1)!} = \frac{1}{3}$$

On simplifying we get

$$\Rightarrow \frac{(r-1)}{(n-r+2)} = \frac{1}{3}$$

 \Rightarrow 3r - 3 = n - r + 2

⇒ n - 4r + 5 =0.....1

Also

 $\frac{\text{the coefficient of } r^{\text{th}} \text{ term}}{\text{coefficient of } r + 1^{\text{th}} \text{ term}} = \frac{3}{5}$

On rearranging we get

 $\xrightarrow{n!}_{(r-1)!(n-r+1)!} \times \frac{r!(n-r)!}{n!} = \frac{3}{5}$

By multiplying

 $\frac{r(r-1)!(n-r)!}{\Rightarrow (r-1)!(n-r+1)!} = \frac{3}{5}$ $\Rightarrow \frac{r(n-r)!}{(n-r+1)!} = \frac{3}{5}$ $\frac{r(n-r)!}{\Rightarrow (n-r+1)(n-r)!} = \frac{3}{5}$

On simplifying we get

 $\frac{r}{\Rightarrow (n-r+1)} = \frac{3}{5}$ Also

 $\frac{\text{the coefficient of } r^{\text{th term}}}{\text{coefficient of } r + 1^{\text{th term}}} = \frac{3}{5}$ $\frac{\frac{n!}{\frac{(r-1)!(n-r+1)!}{n!}}}{\frac{n!}{r!(n-r)!}} = \frac{3}{5}$

On rearranging we get

⇒ 5r = 3n - 3r + 3⇒ 8r - 3n - 3 = 0....2We have 1 and 2 as n - $4r \pm 5 = 0....18r - 3n$ - 3 = 0....2Multiplying equation 1 by number 2 2n - 8r + 10 = 0....3Adding equation 2 and 3 2n - 8r + 10 = 0 -3n - 8r - 3 = 0⇒ -n = -7 n = 7and r = 3



11. Prove that the coefficient of x^n in the expansion of $(1 + x)^{2n}$ is twice the coefficient of x^n in the expansion of $(1 + x)^{2n-1}$.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$

The general term for binomial $(1+x)^{2n}$ is

 $T_{r+1} = {}^{2n}C_r x^r \dots 1$ To find the coefficient of xⁿ r = n T_{n+1} = {}^{2n}C_n x_n The coefficient of xⁿ = ${}^{2n}C_n$ The general term for binomial $(1+x)^{2n-1}$ is T_{r+1} = {}^{2n-1}C_r x_r To find the coefficient of xⁿ Putting n = r T_{r+1} = {}^{2n-1}C_r x_n The coefficient of xⁿ = ${}^{2n-1}C_n$ We have to prove

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

Coefficient of xⁿ in $(1+x)^{2n} = 2$ coefficient of xⁿ in $(1+x)^{2n-1}$ Consider LHS = ${}^{2n}C_n$ = $\frac{2n!}{n! (2n-n)!}$ = $\frac{2n!}{n!}$

Again consider RHS = $2 \times {}^{2n-1}C_n$

$$= 2 \times \frac{(2n-1)!}{n!(2n-1-n)!}$$

$$= 2 \times \frac{(2n-1)!}{n!(n-1)!}$$

Now multiplying and dividing by n we get

$$= 2 \times \frac{(2n-1)!}{n! (n-1)!} \times \frac{n}{n}$$
$$= \frac{2n(2n-1)!}{n! n(n-1)!}$$

$$=\frac{2\pi}{n!n!}$$

From above equations LHS = RHS

Hence the proof.

12. Find a positive value of m for which the coefficient of x^2 in the expansion $(1 + x)^m$ is 6.

Solution:

The general term T_{r+1} in the binomial expansion is given by $T_{r+1} = {}^{n}C_{r} a^{n-r} b^{r}$ Here a = 1, b = x and n = m Putting the value

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

 $T_{r+1} = m C_r 1_{m-r} x_r$ = m Cr XrWe need coefficient of x^2 :. putting r = 2 $T_{2+1} = mC_2 x_2$ The coefficient of $x^2 = {}^mC_2$ Given that coefficient of $x^2 = {}^mC_2 = 6$ $\frac{m!}{2!(m-2)!} = 6$ $\Rightarrow \frac{m(m-1)(m-2)!}{2 \times 1 \times (m-2)!} = 6$ ⇒m $(m - 1) = 12 \Rightarrow m^2 - m - 12$ $=0 \Rightarrow m^2 - 4m + 3m - 12 = 0$ \Rightarrow m (m - 4) + 3 (m - 4) = 0 \Rightarrow (m+3) (m - 4) = 0 \Rightarrow m = - 3, 4 We need positive value of m so m = 4



MISCELLANEOUS EXERCISE

PAGE NO: 175

1. Find a, b and n in the expansion of (a + b)ⁿ if the first three terms of the expansion are 729, 7290 and 30375, respectively.

Solution:

We know that $(r + 1)^{th}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = nC_r a_{n-t} b_r$ The first three terms of the expansion are given as 729, 7290 and 30375 respectively. Then we have, $T_1 = {}^{n}C_0 a^{n-0} b^0 = a^n = 729....1$ $T_2 = {}^{n}C_1 a^{n-1} b^1 = na^{n-1} b = 7290....2$ $T_3 = {}^{n}C_2 a^{n-2} b^2 = n (n-1)/2 a^{n-2} b^2 = 30375.....3$ Dividing 2 by 1 we get $na^{n-1}ba^n = \frac{7290}{720}$ n b a = 10 4 Dividing 3 by 2 we get $n(n-1)a^{n-2}b^22na^{n-1}b = \frac{30375}{7290}$ $\Rightarrow (n-1)b2a = rac{30375}{7290}$ $\Rightarrow (n-1)ba = rac{30375 imes 2}{7290} = rac{25}{3}$ \Rightarrow nba $-\frac{b}{a}=\frac{25}{3}$ $\Rightarrow 10 - ba = \frac{25}{3}$ \Rightarrow ba = 10 - $\frac{25}{3} = \frac{5}{3}$ 5 From 4 and 5 we have n. 5/3 = 10 n = 6Substituting n = 6 in 1 we get a^6 = 729 a = 3

NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

From 5 we have, b/3 = 5/3 b = 5 Thus a = 3, b = 5 and n = 76

2. Find a if the coefficients of x^2 and x^3 in the expansion of $(3 + a x)^9$ are equal.

Solution:

We know that general term of expansion (a + b)ⁿ is

$$T_{r+1} = \left(\frac{n}{r}\right) a^{n-r} b^r$$

For (3+ax)9

EDUGROSS

WISDOMISING KNOWLEDGE

Putting a = 3, $b = a \times \& n = 9$

General term of (3+ax)⁹ is

$$T_{r+1} = \left(\frac{9}{r}\right) 3^{n-r} (ax)^r$$

$$T_{r+1} = \left(\frac{9}{r}\right) 3^{n-r} a^r x^r$$

Since we need to find the coefficients of x^2 and x^3 , therefore

For r = 2

$$T_{2+1} = \left(\frac{9}{2}\right) 3^{n-2} a^2 x^2$$

Thus, the coefficient of $x^2 = \frac{\binom{9}{2}}{3^{n-2}}a^2$

For r = 3

$$T_{3+1} = \left(\frac{9}{3}\right) 3^{n-3} a^3 x^3$$

Thus, the coefficient of $x^3 = \frac{\binom{9}{3}3^{n-3}a^3}{3^n}$ Given that coefficient of x^2 = Coefficient of x^3

$$\Rightarrow \left(\frac{9}{2}\right) 3^{n-2} a^2 = \left(\frac{9}{3}\right) 3^{n-3} a^3$$



$$\Rightarrow \frac{9!}{2!(9-2)!} \times 3^{n-2}a^2 = \frac{9!}{3!(9-3)!} \times 3^{n-3}a^3$$

$$\Rightarrow \frac{3^{n-2}a^2}{3^{n-3}a^3} = \frac{2!(9-2)!}{3!(9-3)!}$$

$$\Rightarrow \frac{3^{(n-2)-(n-3)}}{a} = \frac{2!7!}{3!6!}$$

$$\Rightarrow \frac{3}{a} = \frac{7}{3}$$

$$\therefore a = 9/7$$

Hence, $a = 9/7$

3. Find the coefficient of x^5 in the product $(1 + 2x)^6 (1 - x)^7$ using binomial theorem.

Solution:

 $(1 + 2x)^{6} = {}^{6}C_{0} + {}^{6}C_{1} (2x) + {}^{6}C_{2} (2x)^{2} + {}^{6}C_{3} (2x)^{3} + {}^{6}C_{4} (2x)^{4} + {}^{6}C_{5} (2x)^{5} + {}^{6}C_{6} (2x)^{6}$ = 1 + 6 (2x) + 15 (2x)² + 20 (2x)³ + 15 (2x)⁴ + 6 (2x)⁵ + (2x)⁶ = 1 + 12 x + 60x² + 160 x³ + 240 x⁴ + 192 x⁵ + 64x⁶ (1 - x)⁷ = {}^{7}C_{0} - {}^{7}C_{1} (x) + {}^{7}C_{2} (x)^{2} - {}^{7}C_{3} (x)^{3} + {}^{7}C_{4} (x)^{4} - {}^{7}C_{5} (x)^{5} + {}^{7}C_{6} (x)^{6} - {}^{7}C_{7} (x)^{7} = 1 - 7x + 21x² - 35x³ + 35x⁴ - 21x⁵ + 7x⁶ - x⁷ (1 + 2x)⁶ (1 - x)⁷ = (1 + 12 x + 60x² + 160 x³ + 240 x⁴ + 192 x⁵ + 64x⁶) (1 - 7x + 21x² - 35x³ + 35x⁴ - 21x⁵ + 7x⁶ - x⁷) 192 - 21 = 171 Thus, the coefficient of x⁵ in the correspondent (1 + 2x)⁶ (1 - x)⁷ is 171

Thus, the coefficient of x^5 in the expression $(1+2x)^6(1-x)7$ is 171.

4. If a and b are distinct integers, prove that a – b is a factor of aⁿ – bⁿ, whenever n is a positive integer. [Hint write aⁿ = (a – b + b)ⁿ and expand]

Solution:

In order to prove that (a - b) is a factor of $(a^n - b^n)$, it has to be proved that $a^n - b^n = k (a - b)$ where k is some natural number. a can be written as a = a $-b + b a^n = (a - b + b)^n = [(a - b) + b]^n$ $= {}^nC_0 (a - b)^n + {}^nC_1 (a - b)^{n-1} b + + {}^nC_n b^n$

 $a^{n} - b^{n} = (a - b) [(a - b)^{n-1} + {}^{n}C_{1} (a - b)^{n-1} b + + {}^{n}C_{n} b^{n}]$ $a^{n} - b^{n} = (a - b) k$ Where $k = [(a - b)^{n-1} + {}^{n}C_{1} (a - b)^{n-1} b + + {}^{n}C_{n} b^{n}]$ is a natural number This shows that (a - b) is a factor of $(a^{n} - b^{n})$, where n is positive integer.

5. Evaluate

 $(\sqrt{3}+\sqrt{2})^6-(\sqrt{3}-\sqrt{2})^6$

Solution:

Using binomial theorem the expression $(a + b)^6$ and $(a - b)^6$, can be expanded $(a + b)^6 = {}^6C_0 a^6 + {}^6C_1 a^5 b + {}^6C_2 a^4 b^2 + {}^6C_3 a^3 b^3 + {}^6C_4 a^2 b^4 + {}^6C_5 a b^5 + {}^6C_6 b^6$ $(a - b)^6 = {}^6C_0 a^6 - {}^6C_1 a^5 b + {}^6C_2 a^4 b^2 - {}^6C_3 a^3 b^3 + {}^6C_4 a^2 b^4 - {}^6C_5 a b^5 + {}^6C_6 b^6$ Now $(a + b)^6 - (a - b)^6 = {}^6C_0 a^6 + {}^6C_1 a^5 b + {}^6C_2 a^4 b^2 + {}^6C_3 a^3 b^3 + {}^6C_4 a^2 b^4 + {}^6C_5 a b^5 + {}^6C_6 b^6$ $- [{}^6C_0 a^6 - {}^6C_1 a^5 b + {}^6C_2 a^4 b^2 - {}^6C_3 a^3 b^3 + {}^6C_4 a^2 b^4 - {}^6C_5 a b^5 + {}^6C_6 b^6]$ Now by substituting a = $\sqrt{3}$ and b = $\sqrt{2}$ we get $(\sqrt{3} + \sqrt{2})^6 - (\sqrt{3} - \sqrt{2})^6 = 2 [6 (\sqrt{3})^5 (\sqrt{2}) + 20 (\sqrt{3})^3 (\sqrt{2})^3 + 6 (\sqrt{3}) (\sqrt{2})^5]$ $= 2 [54(\sqrt{6}) + 120 (\sqrt{6}) + 24 \sqrt{6}]$ $= 396 \sqrt{6}$

6. Find the value of

$$\left(a^{2}+\sqrt{a^{2}-1}\right)^{4}+\left(a^{2}-\sqrt{a^{2}-1}\right)^{4}$$

Solution:

Firstly the expression $(x + y)^4 + (x - y)^4$ is simplified by using binomial theorem

$$\begin{aligned} (x+y)^4 &= {}^4 C_0 x^4 + {}^4 C_1 x^3 y + {}^4 C_2 x^2 y^2 + {}^+ C_3 x y^3 + {}^4 C_4 y^4 \\ &= x^4 + 4x^3 y + 6x^2 y^2 + 4xy^3 + y^4 \\ (x-y)^4 &= {}^4 C_0 x^4 - {}^4 C_1 x^3 y + {}^4 C_2 x^2 y^2 - {}^4 C_3 x y^3 + {}^4 C_4 y^4 \\ &= x^4 - 4x^3 y + 6x^2 y^2 - 4xy^3 + y^4 \\ \therefore (x+y)^4 + (x-y)^4 &= 2 (x^4 + 6x^2 y^2 + y^4) \\ \text{Putting } x &= a^2 \text{ and } y = \sqrt{a^2 - 1}, \text{ we obtain} \\ \left(a^2 + \sqrt{a^2 - 1}\right)^4 + \left(a^2 - \sqrt{a^2 - 1}\right)^4 \\ &= 2 \left[(a^2)^4 + 6(a^2)^2 \left(\sqrt{a^2 - 1}\right)^2 + \left(\sqrt{a^2 - 1}\right)^4 \right] \\ &= 2 \left[a^8 + 6a^4 (a^2 - 1) + (a^2 - 1)^2\right] \\ &= 2 \left[a^8 + 6a^6 - 6a^4 + a^4 - 2a^2 + 1\right] \end{aligned}$$

$$=2\left[a^8+6a^6-5a^4-2a^2+1
ight] =2a^8+12a^6-10a^4-4a^2+2$$

7. Find an approximation of $(0.99)^5$ using the first three terms of its expansion.

Solution:

0.99 can be written as 0.99 = 1 - 0.01Now by applying binomial theorem we get (0.99)⁵ = $(1 - 0.01)^5$ = ${}^{5}C_{0}(1)^{5} - {}^{5}C_{1}(1)^{4}(0.01) + {}^{5}C_{2}(1)^{3}(0.01)^{2}$ = $1 - 5(0.01) + 10(0.01)^{2}$ = 1 - 0.05 + 0.001



NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

= 0.951

8. Find n, if the ratio of the fifth term from the beginning to the fifth term from the

end in the expansion of $\left(\sqrt[4]{2} + \frac{1}{\sqrt[4]{3}}\right)^n$ is V6: 1

Solution:

In the expansion $(a + b)^n$, if n is even then the middle term is $(n/2 + 1)^{th}$ term

$${}^{n}C_{4}(\sqrt[4]{2})^{n-1}\left(\frac{1}{\sqrt[4]{3}}\right)^{4} = {}^{n}C_{4}\frac{(\sqrt[4]{2})^{n}}{(\sqrt[4]{2})^{4}} \cdot \frac{1}{3} = {}^{n}C_{4}\frac{(\sqrt[4]{2})^{n}}{2} \cdot \frac{1}{3} = \frac{n!}{6.4!(n-4)!}(\sqrt[4]{2})^{n}$$

$${}^{n}C_{n-4}(\sqrt[4]{2})^{4}\left(\frac{1}{\sqrt[4]{3}}\right)^{n-4} = {}^{n}C_{n-1} \cdot 2 \cdot \frac{(\sqrt[4]{3})^{4}}{(\sqrt[4]{3})^{n}} = {}^{n}C_{n-1} \cdot 2 \cdot \frac{3}{(\sqrt[4]{3})^{n}} = \frac{6n!}{(n-4)!4!} \cdot \frac{1}{(\sqrt[4]{3})^{n}}$$

$$\frac{n!}{6.4!(n-4)!}(\sqrt[4]{2})^{n} : \frac{6n!}{(n-4)!4!} \cdot \frac{1}{(\sqrt[4]{3})^{n}} = \sqrt{6} : 1$$

$$\Rightarrow \frac{(\sqrt[4]{2})^{n}}{6} : \frac{6}{(\sqrt[4]{3})^{n}} = \sqrt{6}$$

$$\Rightarrow (\sqrt[4]{2})^{n} = 36\sqrt{6}$$

$$\Rightarrow (\sqrt[4]{6})^{n} = 36\sqrt{6}$$

$$\Rightarrow 6^{\frac{n}{4}} = 6^{\frac{5}{2}}$$

$$\Rightarrow n = 4 \times \frac{5}{2} = 10$$
Thus the value of n = 10

9. Expand using Binomial Theorem

$$\left(1+\frac{x}{2}-\frac{2}{x}\right)^4, x\neq 0$$



NCERT Solutions for Class 11Maths Chapter 8 Binomial Theorem

Solution:

Using binomial theorem the given expression can be expanded as

Again by using binomial theorem to expand the above terms we get

$$\begin{split} \left(1+\frac{x}{2}\right)^4 &= {}^4C_0\left(1\right)^4 + {}^4C_1\left(1\right)^3\left(\frac{x}{2}\right) + {}^4C_2\left(1\right)^2\left(\frac{x}{2}\right)^2 + {}^4C_3\left(1\right)^1\left(\frac{x}{2}\right)^3 + {}^4C_4\left(\frac{x}{2}\right)^4 \\ &= 1+4\times\frac{x}{2}+6\times\frac{x^2}{4}+4\times\frac{x^3}{8}+\frac{x^4}{16} \\ &= 1+2x+\frac{3x^2}{2}+\frac{x^3}{2}+\frac{x^4}{16} \qquad \dots(2) \\ \left(1+\frac{x}{2}\right)^3 &= {}^3C_0\left(1\right)^3 + {}^3C_1\left(1\right)^2\left(\frac{x}{2}\right) + {}^3C_2\left(1\right)\left(\frac{x}{2}\right)^2 + {}^3C_3\left(\frac{x}{2}\right)^3 \\ &= 1+\frac{3x}{2}+\frac{3x^2}{4}+\frac{x^3}{8} \qquad \dots(3) \end{split}$$

From equation 1, 2 and 3 we get

$$\begin{split} & \left[\left(1 + \frac{x}{2} \right) - \frac{2}{x} \right]^4 \\ &= 1 + 2x + \frac{3x^2}{2} + \frac{x^3}{2} + \frac{x^4}{16} - \frac{8}{x} \left(1 + \frac{3x}{2} + \frac{3x^2}{4} + \frac{x^3}{8} \right) + \frac{8}{x^2} + \frac{24}{x} + 6 - \frac{32}{x^3} + \frac{16}{x^4} \\ &= 1 + 2x + \frac{3}{2}x^2 + \frac{x^3}{2} + \frac{x^4}{16} - \frac{8}{x} - 12 - 6x - x^2 + \frac{8}{x^2} + \frac{24}{x} + 6 - \frac{32}{x^3} + \frac{16}{x^4} \\ &= \frac{16}{x} + \frac{8}{x^2} - \frac{32}{x^3} + \frac{16}{x^4} - 4x + \frac{x^2}{2} + \frac{x^3}{2} + \frac{x^4}{16} - 5 \end{split}$$



10. Find the expansion of $(3x^2 - 2ax + 3a^2)^3$ using binomial theorem.

Solution:

We know that
$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

Putting $a = 3x^2 \& b = -a (2x-3a)$, we get
 $[3x^2 + (-a (2x-3a))]^3$
 $= (3x^2)^3 + 3(3x^2)^2(-a (2x-3a)) + 3(3x^2) (-a (2x-3a))^2 + (-a (2x-3a))^3$
 $= 27x^6 - 27ax^4 (2x-3a) + 9a^2x^2 (2x-3a)^2 - a^3 (2x-3a)^3$
 $= 27x^6 - 54ax^5 + 81a^2x^4 + 9a^2x^2 (4x^2 - 12ax + 9a^2) - a^3 [(2x)^3 - (3a)^3 - 3(2x)^2 (3a) + 3(2x) (3a)^2]$
 $= 27x^6 - 54ax^5 + 81a^2x^4 + 36a^2x^4 - 108a^3x^3 + 81a^4x^2 - 8a^3x^3 + 27a^6 + 36a^4x^2 - 54a^5x$
 $= 27x^6 - 54ax^5 + 117a^2x^4 - 116a^3x^3 + 117a^4x^2 - 54a^5x + 27a^6$
Thus, $(3x^2 - 2ax + 3a^2)^3$
 $= 27x^6 - 54ax^5 + 117a^2x^4 - 116a^3x^3 + 117a^4x^2 - 54a^5x + 27a^6$