

NCERT Solutions for Class 11 Chemistry : Chapter 7 (Equilibrium)

Q.1. A liquid is in equilibrium with its vapour in a sealed container at a fixed temperature. The volume of the container is suddenly increased.

I) What will be the final vapour pressure and what will happen when equilibrium is restored finally?

II) Write down, how initially the rates of evaporation and condensation got changed?

III) Write down the effect observed when there was a change in vapour pressure.


Ans.

(I) Finally, equilibrium will be restored when the rates of the forward and backward processes become equal. However, the vapour pressure will remain unchanged because it depends upon the temperature and not upon the volume of the container.

(II) On increasing the volume of the container, the rates of evaporation will increase initially because now more space is available. Since the amount of the vapours per unit volume decrease on increasing the volume, therefore, the rate of condensation will decrease initially.


(III) On increasing the volume of the container, the vapour pressure will initially decrease because the same amount of vapours are now distributed over a large space.

Q.2. What is K_c for the following equilibrium when the equilibrium concentration of each substance is: $[SO_2] = 0.60M$, $[O_2] = 0.82M$ and $[SO_3] = 1.90M$?



Ans.

As per the question,




$$K_c = \frac{[SO_3]^2}{[SO_2]^2[O_2]}$$

$$= \frac{(1.9)^2 M^2}{(0.6)^2 (0.82) M^3}$$

$$= 12.229 M^{-1} \text{ (approximately)}$$

Hence, K for the equilibrium is $12.229 M^{-1}$.

Q.3. At a certain temperature and total pressure of 105Pa, iodine vapour contains 40% by volume of I atoms



Calculate K_p for the equilibrium

Ans.

Partial pressure of Iodine atoms (I)

$$\begin{aligned}
 p_I &= \frac{40}{100} \times p_{total} \\
 &= \frac{40}{100} \times 10^5 \\
 &= 4 \times 10^4 Pa
 \end{aligned}$$


Partial pressure of I_2 molecules,

$$\begin{aligned}
 p_I &= \frac{60}{100} \times p_{total} \\
 &= \frac{60}{100} \times 10^5 \\
 &= 6 \times 10^4 Pa
 \end{aligned}$$

Now, for the given reaction,

$$\begin{aligned}
 K_p &= \frac{(p_I)^2}{p_{I_2}} = \frac{(4 \times 10^4)^2 Pa^2}{6 \times 10^4 Pa} \\
 &= 2.67 \times 10^4 Pa
 \end{aligned}$$

Q.4. Write the expression for the equilibrium constant, K_c for each of the following reactions:



Ans.

$$K_C = \frac{[NO(g)]^2 [Cl_2(g)]}{[NOCl(g)]^2}$$

$$(ii) K_C = \frac{[CuO(s)]^2 [NO_2(g)]^4 [O_2(g)]}{[Cu(NO_3)_2(g)]^2}$$


$$= [NO_2(g)]^4 [O_2(g)]$$

$$\begin{aligned} \text{(iii)} K_C &= \frac{CH_3COOH_{(aq)}[C_2H_5OH_{(aq)}]}{[CH_3COOC_2H_5_{(aq)}][H_2O_{(l)}]} \\ &= \frac{CH_3COOH_{(aq)}[C_2H_5OH_{(aq)}]}{[CH_3COOC_2H_5_{(aq)}]} \end{aligned}$$

$$\begin{aligned} \text{(iv)} K_C &= \frac{Fe(OH)_{3(s)}}{[Fe^{3+}_{(aq)}][OH^-_{(aq)}]^3} \\ &= \frac{1}{[Fe^{3+}_{(aq)}][OH^-_{(aq)}]^3} \end{aligned}$$

$$\begin{aligned} \text{(v)} K_C &= \frac{[IF_5]^2}{[I_{2(s)}][F_2]^5} \\ &= \frac{[IF_5]^2}{[F_2]^5} \end{aligned}$$

Q.5. Find out the value of K_c for each of the following equilibria from the value of K_p :



Ans.

The relation between K_p and K_c is given as:

$$K_p = K_c(RT)^{\Delta n}$$

(a) Given,

$$R = 0.0831 \text{ barLmol}^{-1}\text{K}^{-1}$$

$$\Delta n = 3 - 2 = 1$$

$$T = 500 \text{ K}$$

$$K_p = 1.8 \times 10^{-2}$$

Now,

$$K_p = K_c(RT)^{\Delta n}$$

$$\Rightarrow 1.8 \times 10^{-2} = K_c(0.0831 \times 500)^1$$

$$\Rightarrow K_c = \frac{1.8 \times 10^{-2}}{0.0831 \times 500}$$

$$= 4.33 \times 10^{-4} (\text{approximately})$$

(b) Here,

$$\Delta n = 2 - 1 = 1$$

$$R = 0.0831 \text{ barLmol}^{-1}\text{K}^{-1}$$

$$T = 1073 \text{ K}$$

$$K_p = 167$$

Now,

$$K_p = K_c (RT)^{\Delta n}$$

$$\Rightarrow 167 = K_c (0.0831 \times 1073)^{\Delta n}$$

$$\Rightarrow K_c = \frac{167}{0.0831 \times 1073}$$

$$= 1.87 (\text{approximately})$$

Q.6. For the following equilibrium, $K_c = 6.3 \times 10^{14}$ at 1000K



Both the forward and reverse reactions in the equilibrium are elementary bimolecular reactions. What is K_c , for the reverse reaction?

Ans.

$$\text{For the reverse reaction, } K_c = \frac{1}{K_c}$$

$$= \frac{1}{6.3 \times 10^{14}}$$

$$= 1.59 \times 10^{-15}$$

Q.7. Explain why solids and pure liquids can be ignored while writing the equilibrium constant expression?

Ans.

This is because molar concentration of a pure solid or liquid is independent of the amount present.

$$\text{Mole concentration} = \frac{\text{Number of moles}}{\text{Volume}}$$

$$\frac{\text{Mass/molecular mass}}{\text{Volume}}$$

$$= \frac{\text{Mass}}{\text{Volume} \times \text{Molecular mass}}$$


$$= \frac{\text{Density}}{\text{Molecular mass}}$$

Though density of solid and pure liquid is fixed and molar mass is also fixed .

∴ Molar concentration are constant.

Q.8.

Reaction between N₂ and O₂ takes place as follows:



If a solution of 0.933 mol of oxygen and 0.482 mol of nitrogen is placed in a 10 L reaction vessel and allowed to form N₂O at a temperature for which $K_c = 2.0 \times 10^{-37}$, determine the composition of equilibrium solution.

Ans.

Let the concentration of N₂O at equilibrium be x.

The given reaction is:



$$[N_2] = \frac{0.482-x}{10}, [O_2] = \frac{0.933-x}{10}, [N_2O] = \frac{x}{10}$$

The value of equilibrium constant is extremely small. This means that only small amounts . Then,

$$[N_2] = \frac{0.482}{10} = 0.0482 \text{ molL}^{-1} \text{ and } [O_2] = \frac{0.933}{10} = 0.0933 \text{ molL}^{-1}$$

Now,

$$K_c = \frac{[N_2O(g)]^2}{[N_2(g)][O_2(g)]}$$

$$\Rightarrow 2.0 \times 10^{-37} = \frac{\left(\frac{x}{10}\right)^2}{(0.0482)^2(0.0933)}$$

$$\Rightarrow \frac{x^2}{100} = 2.0 \times 10^{-37} \times (0.0482)^2 \times (0.0933)$$


$$\Rightarrow x^2 = 43.35 \times 10^{-40}$$

$$\Rightarrow x = 6.6 \times 10^{-20} \quad [N_2O] = \frac{x}{10} = \frac{6.6 \times 10^{-20}}{10}$$

$$= 6.6 \times 10^{-21}$$

Q.9.

Nitric oxide reacts with Br₂ and gives nitrosyl bromide as per reaction given below:



When 0.087 mol of NO and 0.0437 mol of Br₂ are mixed in a closed container at a constant temperature, 0.0518 mol of NOBr is obtained at equilibrium. Calculate equilibrium amount of NO and Br₂.

Ans.

The given reaction is:



Now, 2 mol of NOBr are formed from 2 mol of NO. Therefore, 0.0518 mol of NOBr are formed from 0.0518 mol of NO.

Again, 2 mol of NOBr are formed from 1 mol of Br.

Therefore, 0.0518 mol of NOBr are formed from $\frac{0.0518}{2}$ mol of Br, or 0.0259 mol of NO.

The amount of NO and Br present initially is as follows:

$$[\text{NO}] = 0.087 \text{ mol} \quad [\text{Br}_2] = 0.0437 \text{ mol}$$


Therefore, the amount of NO present at equilibrium is:

$$[\text{NO}] = 0.087 - 0.0518 = 0.0352 \text{ mol}$$

And, the amount of Br present at equilibrium is:

$$[\text{Br}_2] = 0.0437 - 0.0259 = 0.0178 \text{ mol}$$

Q.10. At 450 K, $K_p = 2.0 \times 10^{10}$ /bar for the given reaction at equilibrium.



What is K_c at this temperature?

Ans.)

For the given reaction,

$$\Delta n = 2 - 3 = -1$$

$$T = 450 \text{ K}$$

$$R = 0.0831 \text{ bar L bar K}^{-1} \text{ mol}^{-1}$$

$$K_p = 2.0 \times 10^{10} \text{ bar}^{-1}$$

We know that,

$$K_p = K_c(RT)^{\Delta n}$$

$$\Rightarrow 2.0 \times 10^{10} \text{ bar}^{-1} = K_c(0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 450 \text{ K})^{-1}$$

$$\Rightarrow K_c = \frac{2.0 \times 10^{10} \text{ bar}^{-1}}{(0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 450 \text{ K})^{-1}}$$


$$= (2.0 \times 10^{10} \text{ bar}^{-1})(0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1} \times 450 \text{ K})$$

$$= 74.79 \times 10^{10} \text{ L mol}^{-1}$$

$$= 7.48 \times 10^{11} \text{ L mol}^{-1}$$

$$= 7.48 \times 10^{11} \text{ M}^{-1}$$

Q.11. A sample of HI(g) is placed in flask at a pressure of 0.2 atm. At equilibrium the partial pressure of HI(g) is 0.04 atm. What is K_p for the given equilibrium?



Ans.

The initial concentration of HI is 0.2 atm. At equilibrium, it has a partial pressure of 0.04 atm.

Therefore, a decrease in the pressure of HI is $0.2 - 0.04 = 0.16$. The given reaction is:

2HI(g)		H ₂ (g)	+	I ₂ (g)	
Initial conc.	0.2 atm			0	0
At equilibrium	0.4 atm			0.16	0.16
2	2				
=0.08atm	=0.08atm				

Therefore,

$$K_p = \frac{p_{\text{H}_2} \times p_{\text{I}_2}}{p_{\text{HI}}^2}$$

$$= \frac{0.08 \times 0.08}{(0.04)^2}$$

$$= \frac{0.0064}{0.0016}$$

$$= 4.0$$

Hence, the value of K_p for the given equilibrium is 4.0.


Q.12. A mixture of 1.57 mol of N₂, 1.92 mol of H₂ and 8.13 mol of NH₃ is introduced into a 20 L reaction vessel at 500 K. At this temperature, the equilibrium constant, K_c for the reaction



Is the reaction mixture at equilibrium? If not, what is the direction of the net reaction?

Ans.

The given reaction is:



The given concentration of various species is

$$[N_2] = \frac{1.57}{20} \text{ mol L}^{-1} \quad [H_2] = \frac{1.92}{20} \text{ mol L}^{-1} \quad [NH_3] = \frac{8.31}{20} \text{ mol L}^{-1}$$

Now, reaction quotient Q_c is:

$$Q = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

$$= \frac{\left(\frac{8.31}{20}\right)^2}{\left(\frac{1.57}{20}\right)\left(\frac{1.92}{20}\right)^3}$$

$$= 2.4 \times 10^3$$

Since, $Q \neq K_c$, the reaction mixture is not at equilibrium.

Again, $Q > K_c$. Hence, the reaction will proceed in the reverse direction.


Q.13. The equilibrium constant expression for a gas reaction is,

$$K_c = \frac{[NH_3]^4 [O_2]^5}{[NO]^4 [H_2O]^6}$$


Write the balanced chemical equation corresponding to this expression.

Ans.

The balanced chemical equation corresponding to the given expression can be written as:




Q.14. One mole of H_2O and one mole of CO are taken in 10 L vessel and heated to 725 K. At equilibrium 60% of water (by mass) reacts with CO according to the equation,



Calculate the equilibrium constant for the reaction.

Ans.

The given reaction is:




Compound	H ₂ O	CO	H ₂	CO ₂
Initial Conc.	0.1M	0.1M	0	0
Equilibrium Conc.	0.06M	0.06M	0.04M	0.04M

Therefore, the equilibrium constant for the reaction,

$$K_c = \frac{[H_2][CO_2]}{[H_2O][CO]} = \frac{(0.4 \times 0.4)}{(0.6 \times 0.6)} = 0.444$$

Q.15. At 700 K, equilibrium constant for the reaction



is 54.8. If 0.5 mol L^{-1} of $HI(g)$ is present at equilibrium at 700 K, what are the concentration of $H_2(g)$ and $I_2(g)$ assuming that we initially started with $HI(g)$ and allowed it to reach equilibrium at 700 K?

Ans.

It is given that equilibrium constant K_c for the reaction



Therefore, at equilibrium, the equilibrium constant K'_c for the reaction



$[\text{HI}] = 0.5 \text{ mol L}^{-1}$ will be $1/54.8$.

Let the concentrations of hydrogen and iodine at equilibrium be $x \text{ mol L}^{-1}$

$$[\text{H}_2] = [\text{I}_2] = x \text{ mol L}^{-1}$$

$$\text{Therefore, } \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2} = K_c'$$

$$\Rightarrow \frac{x \times x}{(0.5)^2} = \frac{1}{54.8}$$

$$\Rightarrow x^2 = \frac{0.25}{54.8}$$

$$\Rightarrow x = 0.06754$$

$$x = 0.068 \text{ mol L}^{-1} \text{ (approximately)}$$

Hence, at equilibrium, $[\text{H}_2] = [\text{I}_2] = 0.068 \text{ mol L}^{-1}$.

Q.16. What is the equilibrium concentration of each of the substances in the equilibrium when the initial concentration of ICl was 0.78 M?



Ans.

The given reaction is:

$2\text{ICl}(\text{g})$	\rightleftharpoons	$\text{I}_2(\text{g})$	$+$	$\text{Cl}_2(\text{g})$	
Initial conc.		0.78 M		0	0
At equilibrium		$(0.78 - 2x) \text{ M}$		$x \text{ M}$	$x \text{ M}$

Now, we can write, $\frac{[\text{I}_2][\text{Cl}_2]}{[\text{ICl}]^2} = K_c$

$$\Rightarrow \frac{x \times x}{(0.78 - 2x)^2} = 0.14$$

$$\Rightarrow \frac{x^2}{(0.78 - 2x)^2} = 0.14$$

$$\Rightarrow \frac{x}{0.78 - 2x} = 0.374$$

$$\Rightarrow x = 0.292 - 0.748x$$

$$\Rightarrow 1.748x = 0.292$$

$$\Rightarrow x = 0.167$$


Hence, at equilibrium,

$$[H_2]=[I_2]=0.167 \text{ M}$$

$$[HI]= (0.78 - 2 \times 0.167)M$$

$$= 0.446M$$

Q.17. $K_p = 0.04$ atm at 899 K for the equilibrium shown below. What is the equilibrium concentration of C_2H_6 when it is placed in a flask at 4.0 atm pressure and allowed to come to equilibrium?



Ans.

Let p be the pressure exerted by ethene and hydrogen gas (each) at equilibrium. Now, according to the reaction,

$C_2H_6(g)$		$C_2H_4(g)$	+	$H_2(g)$	
Initial conc.	4.0 M			0	0
At equilibrium	$(4.0-p)$			p	p

We can write,

$$\frac{p_{C_2H_4} \times p_{H_2}}{p_{C_2H_6}} = K_P$$

$$\Rightarrow \frac{p \times p}{4.0-p} = 0.04$$

$$\Rightarrow p^2 = 0.16 - 0.04p$$

$$\Rightarrow p^2 + 0.04p - 0.16 = 0$$

Now,

$$p = \frac{-0.04 \pm \sqrt{(0.04)^2 - 4 \times 1 \times (-0.16)}}{2 \times 1}$$

$$= \frac{-0.04 \pm 0.80}{2}$$

$$= \frac{0.76}{2} \quad (\text{Taking positive value})$$


$$= 0.38$$

Hence, at equilibrium,

$$[C_2H_6] - 4 - p = 4 - 0.38$$

$$= 3.62 \text{ atm}$$

Q.18. Ethyl acetate is formed by the reaction between ethanol and acetic acid and the equilibrium is represented as:



(i) Write the concentration ratio (reaction quotient), Q_c , for this reaction (note: water is not in excess and is not a solvent in this reaction)

(ii) At 293 K, if one starts with 1.00 mol of acetic acid and 0.18 mol of ethanol, there is 0.171 mol of ethyl acetate in the final equilibrium mixture. Calculate the equilibrium constant.

(iii) Starting with 0.5 mol of ethanol and 1.0 mol of acetic acid and maintaining it at 293 K, 0.214 mol of ethyl acetate is found after sometime. Has equilibrium been reached?

Ans.

(i) Reaction quotient,

$$Q_c = \frac{[\text{CH}_3\text{COOC}_2\text{H}_5][\text{H}_2\text{O}]}{[\text{CH}_3\text{COOH}][\text{C}_2\text{H}_5\text{OH}]}$$

(ii) Let the volume of the reaction mixture be V. Also, here we will consider that water is a solvent and is present in excess.

The given reaction is:



Initial conc. 0 0

At equilibrium

= =

Therefore, equilibrium constant for the given reaction is:

$$K_c = \frac{[\text{CH}_3\text{COOC}_2\text{H}_5][\text{H}_2\text{O}]}{[\text{CH}_3\text{COOH}][\text{C}_2\text{H}_5\text{OH}]}$$

$$= \frac{\frac{0.171}{V} \times \frac{0.171}{V}}{\frac{0.829}{V} \times \frac{0.009}{V}} = 3.919$$

$$= 3.92(\text{approximately})$$

(iii) Let the volume of the reaction mixture be V.



Initial conc. 0 0

At equilibrium

= =

Therefore, the reaction quotient is,

$$K_c = \frac{[\text{CH}_3\text{COOC}_2\text{H}_5][\text{H}_2\text{O}]}{[\text{CH}_3\text{COOH}][\text{C}_2\text{H}_5\text{OH}]}$$

$$= \frac{\frac{0.214}{V} \times \frac{0.214}{V}}{\frac{0.786}{V} \times \frac{0.286}{V}} = 0.2037$$

$$= 0.204(\text{approximately})$$


Since $Q_c < K_c$, equilibrium has not been reached.

Q.19. A sample of pure PCl_5 was introduced into an evacuated vessel at 473 K. After equilibrium was attained, concentration of PCl_5 was found to be $0.5 \times 10^{-1} \text{ mol L}^{-1}$. If value of K_c is 8.3×10^{-3} , what are the concentrations of PCl_3 and Cl_2 at equilibrium?



Ans.

Consider the conc. Of both PCl_3 and Cl_2 at equilibrium be $x \text{ mol L}^{-1}$. The given reaction is:



At equilibrium $0.5 \times 10^{-10} \text{ mol L}^{-1}$ $x \text{ mol L}^{-1}$ $x \text{ mol L}^{-1}$

It is given that the value of equilibrium constant, K_c is $8.3 \times 10^{-10} \text{ mol L}^{-3}$

Now we can write the expression for equilibrium as:

$$\frac{[\text{PCl}_2][\text{Cl}_2]}{[\text{PCl}_3]} = K_c$$

$$\Rightarrow \frac{x \times x}{0.5 \times 10^{-10}} = 8.3 \times 10^{-3}$$

$$\Rightarrow x^2 = 4.15 \times 10^{-4}$$

$$\Rightarrow x = 2.04 \times 10^{-2}$$

$$= 0.0204$$

$$= 0.02(\text{approximately})$$

Therefore, at equilibrium,

$$[\text{PCl}_3] = [\text{Cl}_2] = 0.02 \text{ mol L}^{-1}$$

Q.20. One of the reactions that takes place in producing steel from iron ore is the reduction of iron (II) oxide by carbon monoxide to give iron metal and CO_2 .



What are the equilibrium partial pressures of CO and CO_2 at 1050 K if the initial partial pressures are: $p_{\text{CO}} = 1.4 \text{ atm}$ and $p_{\text{CO}_2} = 0.80 \text{ atm}$?

Ans.

For the given reaction,



$$Q_p = \frac{p_{CO_2}}{p_{CO}}$$

$$= \frac{0.80}{1.4}$$

$$= 0.571$$

Since $Q_p > K_p$, the reaction will proceed in the backward direction.

Therefore, we can say that the pressure of CO will increase while the pressure of CO₂ will decrease.

Now, let the increase in pressure of CO = decrease in pressure of CO₂ be p . Then, we can write,

$$K_p = \frac{p_{CO_2}}{p_{CO}}$$

$$\Rightarrow 0.265 = \frac{0.80-p}{1.4+p}$$

$$\Rightarrow 0.371 + 0.265p = 0.80 - p$$

$$\Rightarrow 1.265p = 0.429$$

$$\Rightarrow p = 0.339 \text{ atm}$$

Therefore, equilibrium partial of CO₂, $p_{CO_2} = 0.80 - 0.339 = 0.461$ atm

And, equilibrium partial pressure of CO, $p_{CO} = 1.4 + 0.339 = 1.739$ atm

Q.21. Equilibrium constant, K_c for the reaction



At a specific time, from the analysis we can conclude that composition of the reaction mixture is, 2.0 mol L⁻¹ H₂, 3.0 mol L⁻¹ N₂ and 0.5 mol L⁻¹ NH₃. Find out whether the reaction is at equilibrium or not? Find in which direction the reaction proceeds to reach equilibrium.

Ans.



So,

$$Q_c = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

$$= \frac{(0.5)^2}{(3.0)(2.0)^3}$$

$$= 0.0104$$

It is given that $K_c = 0.061$

∴ $Q_c \neq K_c$, the reaction is not at equilibrium.

$\therefore Q_c < K_c$, the reaction proceeds in the forward direction to reach at equilibrium.

Q.22. Bromine monochloride (BrCl) decays into bromine and chlorine and reaches the equilibrium:




For which $K_c = 42$ at 600 K.

If initially pure BrCl is present at a concentration of $5.5 \times 10^{-5} \text{ mol L}^{-1}$, what is its molar concentration in the mixture at equilibrium?

Ans.)

Let the amount of bromine and chlorine formed at equilibrium be x . The given reaction is:



Now, we can write,

$$\frac{[\text{Br}_2][\text{Cl}_2]}{[\text{BrCl}]^2} = K_c$$

$$\Rightarrow \frac{x \times x}{(5.5 \times 10^{-5} - 2x)^2} = 42$$

$$\Rightarrow \frac{x}{5.5 \times 10^{-5} - 2x} = 6.48$$

$$\Rightarrow x = 35.64 \times 10^{-5} - 12.96x$$

$$\Rightarrow 13.96x = 35.64 \times 10^{-5}$$

$$\Rightarrow x = \frac{35.64}{13.96} \times 10^{-5} = 2.55 \times 10^{-5}$$

So, at equilibrium


$$[\text{BrCl}] = 5.5 \times 10^{-5} - (2 \times 2.55 \times 10^{-5})$$

$$= 5.5 \times 10^{-5} - 5.1 \times 10^{-5}$$

$$= 0.4 \times 10^{-5}$$

$$= 4.0 \times 10^{-6} \text{ mol L}^{-1}$$

Q.23. At 1127 K and 1 atm pressure, a gaseous mixture of CO and CO₂ in equilibrium with solid carbon has 90.55% CO by mass



Calculate K_c for this reaction at the above temperature

Ans.)

Let us assume that the solution is of 100g in total.

Given, mass of CO = 93.55 g

Now, the mass of CO₂ = (100 - 93.55) = 6.45 g

Now, number of moles of CO, $n_{\text{CO}} = \frac{93.5}{28} = 3.34 \text{ mol}$

Number of moles of CO₂, $n_{\text{CO}_2} = \frac{6.45}{44} = 0.146 \text{ mol}$

Partial pressure of CO,

$$P_{\text{CO}} = \frac{n_{\text{CO}}}{n_{\text{CO}} + n_{\text{CO}_2}} \times p_{\text{total}} = \frac{3.34}{3.34 + 0.146} \times 1 = 0.958 \text{ atm}$$

Partial pressure of CO₂, $P_{\text{CO}_2} = \frac{n_{\text{CO}_2}}{n_{\text{CO}} + n_{\text{CO}_2}} \times p_{\text{total}} = \frac{0.146}{3.34 + 0.146} \times 1 = 0.042 \text{ atm}$

Therefore, $K_p = \frac{[\text{CO}]^2}{[\text{CO}_2]}$

$$= \frac{(0.958)^2}{0.042}$$

$$= 14.19$$

For the given reaction,

$$\Delta n = 2 - 1 = 1$$

We know that,


$$K_p = K_c (RT)^{\Delta n}$$

$$\Rightarrow 14.19 = K_c (0.082 \times 1127)^1$$

$$\Rightarrow K_c = \frac{14.19}{0.082 \times 1127}$$

$$= 0.154 (\text{approximately})$$

Q.24. Calculate a) ΔG° and b) the equilibrium constant for the formation of NO_2 from NO and O_2 at 298K



Where;

$$\Delta_f G^\circ (\text{NO}_2) = 52.0 \text{ kJ/mol}$$

$$\Delta_f G^\circ (\text{NO}) = 87.0 \text{ kJ/mol}$$

$$\Delta_f G^\circ (\text{O}_2) = 0 \text{ kJ/mol}$$

Ans.)

(I) We know that,

$$\Delta G^\circ = RT \log K_c$$

$$\Delta G^\circ = 2.303 RT \log K_c$$

$$K_c = \frac{-35.0 \times 10^{-3}}{-2.303 \times 8.314 \times 298}$$


$$= 6.134$$

$$\therefore K_c = \text{antilog}(6.134)$$

$$= 1.36 \times 10^6$$

Therefore, the equilibrium constant for the given reaction K_c is 1.36×10^6

Q.25. Does the number of moles of reaction products increase, decrease or remain same when each of the following equilibria is subjected to a decrease in pressure by increasing the volume?




Ans.





(I) The number of moles of reaction products will increase. According to Le Chatelier's principle, if pressure is decreased, then the equilibrium shifts in the direction in which the number of moles of gases is more. In the given reaction, the number of moles of gaseous products is more than that of gaseous reactants. Thus, the reaction will proceed in the forward direction. As a result, the number of moles of reaction products will increase.

(II) The number of moles of reaction products will decrease.

(III) The number of moles of reaction products remains the same.

Q.26. Which of the following reactions will get affected by increasing the pressure? Also, mention whether change will cause the reaction to go into forward or backward direction.





edugross.com

Ans.)


When pressure is increased:

The reactions given in (i), (iii), (iv), (v), and (vi) will get affected.

Since, the number of moles of gaseous reactants is more than that of gaseous products; the reaction given in (iv) will proceed in the forward direction

Since, the number of moles of gaseous reactants is less than that of gaseous products, the reactions given in (i), (iii), (v), and (vi) will shift in the backward direction

Q.27. The equilibrium constant for the following reaction is 1.6×10^5 at 1024 K.



Find the equilibrium pressure of all gases if 10.0 bar of HBr is introduced into a sealed container at 1024 K.

Ans.

Given, K_p for the reaction i.e., $\text{H}_2(\text{g}) + \text{Br}_2(\text{g}) \rightleftharpoons 2\text{HBr}(\text{g})$ is 1.6×10^5 .

Therefore, for the reaction $2\text{HBr}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{Br}_2(\text{g})$ the equilibrium constant will be,

$$K'_p = \frac{1}{K_p}$$

$$= \frac{1}{1.6 \times 10^5}$$

$$= 6.25 \times 10^{-6}$$

Now, let p be the pressure of both H_2 and Br_2 at equilibrium.

		$\text{H}_2(\text{g})$	+	$\text{Br}_2(\text{g})$	
Initial conc.	10			0	0
At equilibrium	$10-2p$			p	p

Now, we can write,

$$\frac{p_{\text{HBr}} \times p_2}{p_{\text{H}_2}^2} = K'_p$$

$$\frac{p \times p}{(10-2p)^2} = 6.25 \times 10^{-6}$$

$$\frac{p}{10-2p} = 2.5 \times 10^{-3}$$

$$p = 2.5 \times 10^{-2} - (5.0 \times 10^{-3})p$$

$$p + (5.0 \times 10^{-3})p = 2.5 \times 10^{-2}$$

$$(1005 \times 10^{-3}) = 2.5 \times 10^{-2}$$


$$p = 2.49 \times 10^{-2} \text{ bar} = 2.5 \times 10^{-2} \text{ bar (approximately)}$$

Therefore, at equilibrium,

$$[\text{H}_2] = [\text{Br}_2] = 2.49 \times 10^{-2} \text{ bar} \quad [\text{HBr}] = 10 - 2 \times (2.49 \times 10^{-2}) \text{ bar}$$

$$= 9.95 \text{ bar} = 10 \text{ bar (approximately)}$$

Q.28. Dihydrogen gas is obtained from natural gas by partial oxidation with steam as per following endothermic reaction:



(I) Write an expression for K_p for the above reaction.

(II) How will the values of K_p and composition of equilibrium mixture be affected by

- (i) Increasing the pressure
- (ii) Increasing the temperature
- (iii) Using a catalyst?

Ans.)

(I) For the given reaction,

$$K_p = \frac{p_{\text{CO}} \times p_{\text{H}_2}^3}{p_{\text{CH}_4} \times p_{\text{H}_2\text{O}}}$$

(II) (i) According to Le Chatelier's principle, the equilibrium will shift in the backward direction.

(ii) According to Le Chatelier's principle, as the reaction is endothermic, the equilibrium will shift in the forward direction.

(iii) The equilibrium of the reaction is not affected by the presence of a catalyst. A catalyst only increases the rate of a reaction. Thus, equilibrium will be attained quickly.


Q.29. Describe the effect of:

I) Removal of CO

II) Addition of H_2

III) Removal of CH_3OH on the equilibrium of the reaction:

IV) Addition of CH_3OH



Ans.)

- (I) On removing CO, the equilibrium will shift in the backward direction.
 (II) According to Le Chatelier's principle, on addition of H₂, the equilibrium of the given reaction will shift in the forward direction.
 (III) On removing CH₃OH, the equilibrium will shift in the forward direction.
 (IV) On addition of CH₃OH, the equilibrium will shift in the backward direction.

Q.30. At 473 K, equilibrium constant K_c for decomposition of phosphorus pentachloride, PCl_5 is $8.3 \times$

10^{-3} . If decomposition is depicted as,



$$\Delta_r H^\circ = 124.0 \text{ kJmol}^{-1}$$

- a) Write an expression for K_c for the reaction.
 b) What is the value of K_c for the reverse reaction at the same temperature?
 c) What would be the effect on K_c if
 (i) more PCl_5 is added
 (ii) pressure is increased?
 (iii) The temperature is increased?

Ans.)

$$(a) K_c = \frac{[PCl_3(g)][Cl_2(g)]}{[PCl_5(g)]}$$

(b) Value of K_c for the reverse reaction at the same temperature is:


$$K_c' = \frac{1}{K_c}$$

$$= \frac{1}{8.3 \times 10^{-3}} = 1.2048 \times 10^2$$

$$= 120.48$$

- (c)(i) K_c would remain the same because in this case, the temperature remains the same.
 (ii) K_c is constant at constant temperature. Thus, in this case, K_c would not change. (iii) In an endothermic reaction, the value of K_c increases with an increase in temperature. Since the given reaction is an endothermic reaction, the value of K_c will increase if the temperature is increased.

Q.31. Dihydrogen gas used in Haber's process is produced by reacting methane from natural gas with high temperature steam. The first stage of two stage reaction involves the formation of CO and H₂. In second stage, CO formed in first stage is reacted with more steam in water gas shift reaction,




If a reaction vessel at 400°C is charged with an equimolar mixture of CO and steam such that $P_{CO} = P_{H_2O}$

= 4.0 bar, what will be the partial pressure of H₂ at equilibrium? $K_p = 10.1$ at 400°C

Ans.)

Let the partial pressure of both carbon dioxide and hydrogen gas be p . The given reaction is:



Initial conc. 4.0 bar 4.0 bar 0 0

At equilibrium 4.0-p 4.0-p p p

Given $K_p = 10.1$

$$\frac{P_{\text{CO}_2} \times P_{\text{H}_2}}{P_{\text{CO}} \times P_{\text{H}_2\text{O}}} = K_P$$

$$\Rightarrow \frac{p \times p}{(4.0-p)(4.0-p)} = 10.1$$

$$\Rightarrow \frac{p}{4.0-p} = 3.178$$

$$\Rightarrow p = 12.712 - 3.178p$$


$$4.178p = 12.712$$

$$p = \frac{12.712}{4.178}$$

$$p = 3.04$$

So, partial pressure of H₂ is 3.04 bar at equilibrium.

Q.32. Predict which of the following reaction will have appreciable concentration of reactants and products:



Ans.)


If the value of K_c lies between 10^{-3} and 10^3 , a reaction has appreciable concentration of reactants and products. Thus, the reaction given in (c) will have appreciable concentration of reactants and products.

Q.33. The value of K_c for the reaction $3\text{O}_2\text{(g)} \rightleftharpoons 2\text{O}_3\text{(g)}$ is 2.0×10^{-50} at 25°C. If the equilibrium

concentration of O_2 in air at 25°C is 1.6×10^{-2} , what is the concentration of O_3 ?

Ans.)

Given,



$$\text{Then, } K_c = \frac{[O_3(g)]^2}{[O_2(g)]^3}$$

Given that $K_c = 2.0 \times 10^{-50}$ and $[O_2(g)] = 1.6 \times 10^{-2}$

Then,

$$2.0 \times 10^{-50} = \frac{[O_3(g)]^2}{[1.6 \times 10^{-2}]^3}$$

$$\Rightarrow [O_3(g)]^2 = 2.0 \times 10^{-50} \times (1.6 \times 10^{-2})^3$$

$$\Rightarrow [O_3(g)]^2 = 8.192 \times 10^{-56}$$

$$\Rightarrow [O_3(g)] = 2.86 \times 10^{-28} M$$

So, the conc. of O_3 is $2.86 \times 10^{-28} M$.

Q.34. The reaction, $CO(g) + 3H_2(g) \rightarrow CH_4(g) + H_2O(g)$ at 1300K is at equilibrium in a 1L

container. It has 0.30 mol of CO, 0.10 mol of H_2 and 0.02 mol of H_2O and y amount of CH_4 in the

container. Find the concentration of CH_4 in the mixture.

The equilibrium constant, K_c is 3.90 at the given temp.

Ans.)

Let the concentration of CH_4 at equilibrium be y.



At equilibrium,

$$\text{For CO} - \frac{0.3}{1} = 0.3M$$

$$\text{For } H_2 - \frac{0.1}{1} = 0.1M$$

$$\text{For } H_2O - \frac{0.02}{1} = 0.02M \quad K_c = 3.90$$

Therefore,

$$\frac{[CH_4(g)][H_2O(g)]}{[CO(g)][H_2(g)]^3} = K_c \frac{y \times 0.02}{0.3 \times (0.1)^3} = 3.9 \quad y = \frac{3.9 \times 0.3 \times (0.1)^3}{0.02} \quad y = \frac{0.00117}{0.02} \quad y =$$

$$0.0585M \quad y = 5.85 \times 10^{-2}M$$

Therefore, the concentration of CH_4 at equilibrium is $5.85 \times 10^{-2}M$

Q.35. What is conjugate acid-base pair? Find the conjugate acid/base of the given species:

(i) HNO_2

(ii) CN^-

(iii) $HClO_4$

(iv) F^-

(v) OH^-

(vi) CO_3^{2-}

(vii) S^-

Ans.)



A conjugate acid-base pair is a pair that has a difference of only one proton.

The conjugate acid-base pair of the following are as follows:



(i) $HNO_2 - NO_2^-$ (Base)

(ii) $CN^- - HCN$ (Acid)

(iii) $HClO_4 - ClO_4^-$ (Base)



Q.36. Which of the followings are Lewis acids?



Ans.)

Lewis acids are the acids which can accept a pair of electrons.






Q.37. What will be the conjugate bases for the Brønsted acids:



Ans.)

The following shows the conjugate bases for the Bronsted acids:



Q.38. Write the conjugate acids for the following Brönsted bases:



Ans.)

	Brönsted base	Conjugate acid
1	NH_3	NH_4^+
2	HCOO^-	HCOOH
3	NH_2^-	NH_3

Q.39. The species: H_2O , HCO_3^- , HSO_4^- and NH_3 can act both as Brönsted acids and bases. For each case give the corresponding conjugate acid and base.

Ans.)

	Species	Conjugate base	Conjugate acid
1	HCO_3^-	CO_3^{2-}	H_2CO_3
2	HSO_4^-	SO_4^{2-}	H_2SO_4
3	NH_3	NH_2^-	NH_4^+
4	H_2O	OH^-	H_3O^+

Q.40. Classify the following species into Lewis acids and Lewis bases and show that these species act as Lewis base/acid:



Ans.)



It is a Lewis acid as it has tendency to accept a pair of electrons.



It is a Lewis acid as it has tendency to accept a pair of electrons.



It is a Lewis base as it has tendency to lose a pair of electrons.



It is a Lewis base as it has tendency to lose its lone pair of electrons.

Q.41. The concentration of hydrogen ion in a sample of soft drink is 3.8×10^{-3} M. what is its pH?

Ans.)

$$\text{pH} = -\log[\text{H}^+]$$

$$= -\log(2.5 \times 10^{-4})$$

$$= -\log 2.5 - \log 10^{-4}$$

$$= -\log 2.5 + 4$$

$$= -0.398 + 4$$

$$= 3.602$$

Q.42. The pH of a sample of vinegar is 3.76. Calculate the concentration of hydrogen ion in it.

Ans.)

$$pH = -\log[H^+]$$

$$\Rightarrow \log[H^+] = -pH$$

$$\Rightarrow [H^+] = \text{antilog}(-pH)$$

$$= \text{antilog}(-2.36)$$

$$= 0.004365$$

$$= 4.37 \times 10^{-3} \therefore 4.37 \times 10^{-3} \text{ is the concentration of white vinegar sample.}$$

Q.43. Ionization constant for the following acids are given:

$$\text{HF} = 5.7 \times 10^{-5} \text{ at } 298\text{K}$$

$$\text{HCOOH} = 1.7 \times 10^{-3} \text{ at } 298\text{K}$$

$$\text{HCN} = 3.7 \times 10^{-8} \text{ at } 298\text{K}$$

Find out the conjugate bases for the above acids.

Ans.)

$$\text{For } F^-, K_b = \frac{K_w}{K_a} = \frac{10^{-14}}{(5.7 \times 10^{-5})} = 1.75 \times 10^{-9}$$

$$\text{For } \text{HCOO}^-, K_b = \frac{10^{-14}}{(1.7 \times 10^{-3})} = 5.88 \times 10^{-11}$$

$$\text{For } \text{CN}^-, K_b = \frac{10^{-14}}{(3.7 \times 10^{-8})} = \times 10^{-11} = 2.70 \times 10^{-6}$$

Q.44. The ionization constant of phenol is 1.0×10^{-10} . What is the concentration of phenolate ion in 0.05 M solution of phenol? What will be its degree of ionization if the solution is also 0.01M in sodium phenolate?

Ans.)

$\text{C}_6\text{H}_5\text{OH}$	$\text{C}_6\text{H}_5\text{O}^- + \text{H}^+$		
Initial	0.06M		
After dissociation	$0.06-x$	x	x

$$\therefore K_a = \frac{x \times x}{0.06 - x} = 1.0 \times 10^{-8}$$

$$\Rightarrow \frac{x^2}{0.06} = 1.0 \times 10^{-8}$$

$$\Rightarrow x^2 = 6 \times 10^{-10}$$

$$\Rightarrow x = 2.4 \times 10^{-5} M$$

In presence of 0.02 sodium phenolate(C_6H_5Na), suppose y is the amount of phenol dissociated, then at equilibrium

$$[C_6H_5OH] = 0.06 - y \simeq 0.06,$$

$$[C_6H_5O^-] = 0.02 + y \simeq 0.02M,$$

$$[H^+] = y M$$

$$\therefore K_a = \frac{(0.02)(y)}{0.06} = 1.0 \times 10^{-8}$$

$$\Rightarrow y = \frac{1.0 \times 0.06}{(0.02)} \times 10^{-8}$$

$$\Rightarrow y = 6 \times 10^{-8} \therefore \text{degree of ionization} = \alpha = \frac{y}{c} = \frac{6 \times 10^{-8}}{6 \times 10^{-2}} (\text{Here } c = 0.06 = 6 \times$$

$$10^{-2}) = 10^{-6}$$

$$\text{So, } \alpha = 10^{-6}$$

Q.45 The first ionization constant of H_2S is 9.1×10^{-8} . Calculate the concentration of HS^- ion in its 0.1M solution. How will this concentration be affected if the solution is 0.1M in HCl also? If the second dissociation constant of H_2S is 1.2×10^{-13} , calculate the concentration of S^{2-} under both conditions.

Ans.)

To calculate $[HS^-]$

To find $[HS^-]$:

Case 1 - HCl is absent.

Concentration	H_2S	H^+	HS^-
Initial Conc.	0.1	0	0
Final Conc.	$0.1 - x$	x	x

Now,

$$K_a = \frac{[H^+][HS^-]}{[H_2S]} = 9.1 \times 10^{-8} \text{ (given)}$$

$$\text{Therefore, } \frac{x^2}{(0.1-x)} = 9.1 \times 10^{-8}$$

But $0.1-x$ is approximately equal to 0.1 . Substituting this value in the equation:

$$x^2/0.1 = 9.1 \times 10^{-8}; x^2 = 9.1 \times 10^{-9}; x = 9.54 \times 10^{-5} \text{ M}$$

Therefore, concentration of HS^- is $9.54 \times 10^{-5} \text{ M}$

Case 2 – HCl is present

.

Concentration	H_2S	H^+	HS^-
Initial Conc.	0.1	0	0.1
Final Conc.	$0.1-y$	y	$0.1+y$

Now,

$$K_a = \frac{[\text{H}^+][\text{HS}^-]}{[\text{H}_2\text{S}]} = \frac{(y \cdot (0.1+y))}{(0.1-y)} = 9.1 \times 10^{-8} \text{ (given)}$$

But $(0.1 + y)$ and $(0.1 - y)$ can be approximated to 0.1

Therefore,

$$9.1 \times 10^{-8} = (0.1 \cdot y)/0.1$$

$$\text{Therefore, } y = [\text{HS}^-] = 9.1 \times 10^{-8} \text{ M}$$

To calculate $[\text{S}^{2-}]$:

Case 1 – HCl is absent.

The dissociation of HS^- is given by the equation: $\text{HS}^- \rightleftharpoons \text{H}^+ + \text{S}^{2-}$

$$[\text{HS}^-] = 9.1 \times 10^{-5} \text{ M}; [\text{H}^+] = 9.54 \times 10^{-5} \text{ M}$$

$$K_a = \frac{[\text{H}^+][\text{S}^{2-}]}{[\text{HS}^-]} = 1.2 \times 10^{-13} \text{ (given)}$$

$$K_a = (9.1 \times 10^{-5} \cdot [\text{S}^{2-}])/9.1 \times 10^{-5}$$

$$\text{Therefore, } [\text{S}^{2-}] = 1.2 \times 10^{-13} \text{ M}$$

Case 2 – HCl is present

$$[\text{HS}^-] = 9.1 \times 10^{-8} \text{ M}; [\text{H}^+] = 0.1 \text{ M}$$

$$K_a = 1.2 \times 10^{-13} \text{ M} = (0.1 \cdot [\text{S}^{2-}])/9.1 \times 10^{-8}$$

$$\text{Therefore, } [\text{S}^{2-}] = 1.092 \times 10^{-10} \text{ M}$$

Q.46. The ionization constant of acetic acid is 1.74×10^{-5} . Calculate the degree of dissociation of acetic acid in its 0.05 M solution. Calculate the concentration of acetate ion in the solution and its pH

Ans.)



$$K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]} = \frac{[\text{H}^+]^2}{[\text{CH}_3\text{COOH}]}$$

$$\Rightarrow [\text{H}^+] = \sqrt{K_a[\text{CH}_3\text{COOH}]} = \sqrt{(1.74 \times 10^{-5})(5 \times 10^{-2})} = 9.33 \times 10^{-4} \text{ M}$$

$$[\text{CH}_3\text{COO}^-] = [\text{H}^+] = 9.33 \times 10^{-4} \text{ M}$$

$$\text{pH} = -\log(9.33 \times 10^{-4}) = 4 - 0.9699 = 4 - 0.97 = 3.03$$

Q.47. It has been found that the pH of a 0.01 M solution of an organic acid is 4.15 . Calculate the concentration of the anion, the ionization constant of the acid and its $\text{p}K_a$.

Ans.)



$$\text{pH} = -\log[\text{H}^+]$$

$$\log[\text{H}^+] = -4.15$$

$$[\text{H}^+] = 7.08 \times 10^{-5} \text{ M} \quad [\text{A}^-] = [\text{H}^+] = 7.08 \times 10^{-5} \text{ M} \quad K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} =$$

$$\frac{(7.08 \times 10^{-5})(7.08 \times 10^{-5})}{10^{-2}} = 5.0 \times 10^{-7}$$

$$\text{p}K_a = -\log K_a = -\log(5.0 \times 10^{-7}) = 7 - 0.699 = 6.301$$

Q.48. Assuming complete dissociation, calculate the pH of the following solutions:

(I) 0.004 M HCl

(II) 0.003 M NaOH

(III) 0.002 M HBr

(IV) 0.002 M KOH

Ans.)



$$\therefore [\text{H}^+] = [\text{HCl}] = 4 \times 10^{-3} \text{ M}$$

$$\text{pH} = -\log(4 \times 10^{-3}) = 2.398$$



$$\therefore [\text{OH}^-] = 3 \times 10^{-3} \text{ M}$$

$$[\text{H}^+] = \frac{10^{-14}}{(3 \times 10^{-3})} = 3 \times 10^{-12} \text{ M}$$

$$\text{pH} = -\log(3 \times 10^{-12}) = 11.52$$



$$\therefore [\text{H}^+] = 2 \times 10^{-3} \text{ M}$$

$$\text{pH} = -\log(2 \times 10^{-3} \text{ M}) = 2.70$$



$$\therefore [\text{OH}^-] = 2 \times 10^{-3} \text{ M}$$

$$[\text{H}^+] = \frac{10^{-14}}{2} = 5 \times 10^{-12}$$

$$pH = -\log(5 \times 10^{-12}) = 11.30$$

Q.49. Calculate the pH of the following solutions:

(I) 2g of TlOH dissolved in water to give 2 litre of the solution

(II) 0.3g of Ca(OH)₂ dissolved in water to give 500mL of the solution

(III) 0.3g of NaOH dissolved in water to give 200mL of the solution

(IV) 1 mL of 13.6 M HCl is diluted with water to give 1 litre of the solution

Ans.)

$$(I) \text{Molar conc. of TlOH} = \frac{2g}{(204+16+1)g \text{ mol}^{-1}} \times \frac{1}{2L} = 4.52 \times 10^{-3} M$$

$$[OH^-] = [TlOH] = 4.52 \times 10^{-3} M$$

$$[H^+] = \frac{10^{-14}}{(4.52 \times 10^{-3})} = 2.21 \times 10^{-12} M$$

$$\therefore pH = -\log(2.21 \times 10^{-12}) = 12 - (0.3424) = 11.66$$

$$(II) \text{Molar conc. of Ca(OH)}_2 = \frac{0.3g}{(40+34)g \text{ mol}^{-1}} \times \frac{1}{0.5L} = 8.11 \times 10^{-3} M$$

$$[OH^-] = 2[Ca(OH)_2] = 2 \times (8.11 \times 10^{-3}) M = 16.22 \times 10^{-3} M$$

$$pOH = -\log(16.22 \times 10^{-3}) = 3 - 1.2101 = 1.79$$

$$pH = 14 - 1.79 = 12.21$$

$$(III) \text{Molar conc. of NaOH} = \frac{0.3g}{(40+34)g \text{ mol}^{-1}} \times \frac{1}{0.2L} = 3.75 \times 10^{-2} M$$

$$[OH^-] = 3.75 \times 10^{-2} M$$

$$pOH = -\log(3.75 \times 10^{-2}) = 2 - 0.0574 = 1.43$$

$$pH = 14 - 1.43 = 12.57$$

$$(IV) M_1 V_1 = M_2 V_2$$

$$\therefore 13.6M \times 1mL = M_2 \times 1000mL$$


$$\therefore M_2 = 1.36 \times 10^{-2} M$$

$$[H^+] = [HCl] = 1.36 \times 10^{-2} M$$

$$pH = -\log(1.36 \times 10^{-2}) = 2 - 0.1335 \simeq 1.87$$

Q.50. The degree of ionization of a 0.1M bromoacetic acid solution is 0.132. Calculate the pH of the solution and the pK_a of bromoacetic acid.

Ans.)



$$K_a = \frac{C\alpha \cdot C\alpha}{C(1-\alpha)} = \frac{C\alpha^2}{1-\alpha} \simeq C\alpha^2 = 0.1 \times (0.132)^2 = 1.74 \times 10^{-3}$$


$$pK_a = -\log(1.74 \times 10^{-3}) = 3 - 0.2405 = 2.76$$

$$[H^+] = C\alpha = 0.1 \times 0.132 = 1.32 \times 10^{-2} M$$

$$pH = -\log(1.32 \times 10^{-2}) = 2 - 0.1206 = 1.88$$

Q.51. The degree of ionization of a 0.1M bromoacetic acid solution is 0.132. Calculate the pH of the solution and the pK_a of bromoacetic acid.

Ans.)



$$K_a = \frac{C\alpha \cdot C\alpha}{C(1-\alpha)} = \frac{C\alpha^2}{1-\alpha} \simeq C\alpha^2 = 0.1 \times (0.132)^2 = 1.74 \times 10^{-3}$$

$$pK_a = -\log(1.74 \times 10^{-3}) = 3 - 0.2405 = 2.76$$

$$[H^+] = C\alpha = 0.1 \times 0.132 = 1.32 \times 10^{-2} M$$

$$pH = -\log(1.32 \times 10^{-2}) = 2 - 0.1206 = 1.88$$

Q.52. What is the pH of 0.001 M aniline solution? The ionization constant of aniline can be taken from Table 7.7. Calculate the degree of ionization of aniline in the solution. Also calculate the ionization constant of the conjugate acid of aniline.

Ans.)

$$K_b = 4.27 \times 10^{-10}$$

$$c = 0.001M$$

$$pH = ?$$

$$\alpha = ?$$

$$K_b = c\alpha^2$$

$$4.27 \times 10^{-10} = 0.001 \times \alpha^2$$

$$4270 \times 10^{-10} = \alpha^2$$

$$65.34 \times 10^{-5} = \alpha = 6.53 \times 10^{-4}$$

$$\text{Then, [anion]} = c\alpha = 0.001 \times 65.34 \times 10^{-5} = 0.065 \times 10^{-5} \quad pOH =$$

$$-\log(0.065 \times 10^{-5})$$

$$= 6.187$$

$$pH = 7.813$$

Now,

$$K_a \times K_b = K_w$$

$$K_a = \frac{10^{-14}}{4.27 \times 10^{-10}}$$

$$= 2.34 \times 10^{-5} \therefore 2.34 \times 10^{-5} \text{ is the ionization constant.}$$

Q.53. Calculate the degree of ionization of 0.05M acetic acid if its p_{K_a} value is 4.74. How is the degree of dissociation affected when its solution also contains

(I) 0.01 M

(II) 0.1 M in HCl?

Ans.)

$$c = 0.05M$$

$$p_{K_a} = 4.74$$

$$p_{K_a} = -\log(K_a)$$

$$K_a = c\alpha^2$$

$$\alpha = \sqrt{\frac{K_a}{c}}$$

$$\alpha = \sqrt{\frac{1.82 \times 10^{-5}}{5 \times 10^{-2}}} = 1.908 \times 10^{-2}$$

When HCl is added to the solution, the concentration of H⁺ ions will increase. Therefore, the equilibrium will shift in the backward direction i.e., dissociation of acetic acid will decrease.

Case 1: When 0.01 M HCl is taken.

Let x be the amount of acetic acid dissociated after the addition of HCl.

CH ₃ COOH	H ⁺ + CH ₃ COO ⁻		
Initial conc.	0.05M	0	0
After dissociation	0.05-x	0.01+x	x

As the dissociation of a very small amount of acetic acid will take place, the values i.e., 0.05 - x and 0.01 + x can be taken as 0.05 and 0.01 respectively.

$$K_a = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$$

$$\therefore = \frac{(0.01)x}{0.05}$$

$$x = \frac{1.82 \times 10^{-5} \times 0.05}{0.01}$$

$$x = 1.82 \times 10^{-3} \times 0.05M$$

Now,

$$\alpha = \frac{\text{Amount of acid dissociation}}{\text{Amount of acid taken}}$$

$$= \frac{1.82 \times 10^{-3} \times 0.05}{0.05}$$

$$= 1.82 \times 10^{-3}$$

Case 2: When 0.1 M HCl is taken.

Let the amount of acetic acid dissociated in this case be X. As we have done in the first case, the concentrations of various species involved in the reaction are:

$$[CH_3COOH] = 0.05 - X; 0.05M$$

$$[CH_3COO^-] = X$$

$$[H^+] = 0.1 + X; 0.1M$$

$$K_a = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$$

$$\therefore K_a = \frac{(0.1)x}{0.05}$$

$$x = \frac{1.82 \times 10^{-5} \times 0.05}{0.1}$$

$$x = 1.82 \times 10^{-4} \times 0.05M$$

Now,

$$\alpha = \frac{\text{Amount of acid dissociation}}{\text{Amount of acid taken}}$$

$$= \frac{1.82 \times 10^{-4} \times 0.05}{0.05}$$

$$= 1.82 \times 10^{-4}$$

Q.54. The ionization constant of dimethylamine is 5.4×10^{-4} . Calculate its degree of ionization in its 0.02 M solution. What percentage of dimethylamine is ionized if the solution is also 0.1 M in NaOH?

Ans.)

$$K_b = 5.4 \times 10^{-4}$$

$$c = 0.02M$$

$$\text{Then, } \alpha = \sqrt{\frac{K_b}{c}}$$

$$= \sqrt{\frac{5.4 \times 10^{-4}}{0.02}} = 0.1643$$

Now, if 0.1 M of NaOH is added to the solution, then NaOH (being a strong base) undergoes complete ionization.



$$0.1M \quad 0.1M$$

And,



$$(0.02-x) \qquad \qquad \qquad x \qquad \qquad \qquad x$$

$$;0.02M \qquad \qquad \qquad \qquad \qquad \qquad \qquad ;0.1M$$

$$\text{Then, } [(\text{CH}_3)_2\text{NH}_2^+] = x$$

$$[\text{OH}^-] = x + 0.1; 0.1$$

$$\Rightarrow K_b = \frac{[(\text{CH}_3)_2\text{NH}_2^+][\text{OH}^-]}{[(\text{CH}_3)_2\text{NH}]}$$

$$5.4 \times 10^{-4} = \frac{x \times 0.1}{0.02}$$

$$x = 0.0054$$

It means that in the presence of 0.1 M NaOH, 0.54% of dimethylamine will get dissociated.

Q.55. Calculate the hydrogen ion concentration in the following biological fluids whose pH are given below:

(I) Human saliva, 6.4

(II) Human stomach fluid, 1.2

(III) Human muscle-fluid, 6.83

(IV) Human blood, 7.38

Ans.)

(I) Human saliva, 6.4:

$$\text{pH} = 6.4$$

$$6.4 = -\log [\text{H}^+] \quad [\text{H}^+] = 3.98 \times 10^{-7}$$

(II) Human stomach fluid, 1.2:

$$\text{pH} = 1.2$$

$$1.2 = -\log [\text{H}^+] \quad \therefore [\text{H}^+] = 0.063$$

(III) Human muscle fluid 6.83:

$$\text{pH} = 6.83$$

$$\text{pH} = -\log [\text{H}^+]$$

$$6.83 = -\log [\text{H}^+] \quad [\text{H}^+] = 1.48 \times 10^{-7} \text{ M}$$

(IV) Human blood, 7.38:

$$\text{pH} = 7.38 = -\log [\text{H}^+] \quad [\text{H}^+] = 4.17 \times 10^{-8} \text{ M}$$

Q.56. The pH of milk, black coffee, tomato juice, lemon juice and egg white are 6.8, 5.0, 4.2, 2.2 and 7.8 respectively. Calculate corresponding hydrogen ion concentration in each.

Ans.)

The hydrogen ion concentration in the given substances can be calculated by using the given relation: $\text{pH} = -\log [\text{H}^+]$

$$(I) \text{pH of milk} = 6.8$$

$$\text{Since, } \text{pH} = -\log [\text{H}^+]$$

$$6.8 = -\log [\text{H}^+] \quad \log$$

$$[\text{H}^+] = -6.8$$

$$[\text{H}^+] = \text{antilog}(-6.8)$$

$$= 1.5 \times 10^{-7} M$$

(II) pH of black coffee = 5.0

$$\text{Since, } \text{pH} = -\log [\text{H}^+]$$

$$5.0 = -\log [\text{H}^+] \log$$

$$[\text{H}^+] = -5.0$$

$$[\text{H}^+] = \text{antilog}(-5.0)$$

$$= 10^{-5} M$$

(III) pH of tomato = 4.2

$$\text{Since, } \text{pH} = -\log [\text{H}^+]$$

$$4.2 = -\log [\text{H}^+] \log$$

$$[\text{H}^+] = -4.2$$

$$[\text{H}^+] = \text{antilog}(-4.2)$$

$$= 6.31 \times 10^{-5} M$$

(IV) pH of lemon juice = 2.2

$$\text{Since, } \text{pH} = -\log [\text{H}^+]$$

$$2.2 = -\log [\text{H}^+] \log$$

$$[\text{H}^+] = -2.2$$

$$[\text{H}^+] = \text{antilog}(-2.2)$$

$$= 6.31 \times 10^{-3} M$$

(V) pH of egg white = 7.8

$$\text{Since, } \text{pH} = -\log [\text{H}^+]$$

$$7.8 = -\log [\text{H}^+] \log$$

$$[\text{H}^+] = -7.8$$

$$[\text{H}^+] = \text{antilog}(-7.8)$$

$$= 1.58 \times 10^{-8} M$$

Q.57. If 0.561 g of KOH is dissolved in water to give 200 mL of solution at 298 K. Calculate the concentrations of potassium, hydrogen and hydroxyl ions. What is its pH?

Ans.)

$$[\text{KOH}]_{\text{aq}} = \frac{0.561}{\frac{1}{5}} g/L$$

$$= 2.805 g/L$$

$$= 2.805 \times \frac{1}{56.11} M$$

$$= 0.05 \times 56.11$$

$$= 0.05M$$



$$[OH^-] = 0.05M = [K^+] \quad [H^+][H^-] = K_w$$

$$[H^+] = \frac{K_w}{[OH^-]} = \frac{10^{-14}}{0.05} = 2 \times 10^{-13}M$$

$$\therefore pH = 12.70$$

Q.58. The solubility of $Sr(OH)_2$ at 298 K is 19.23 g/L of solution. Calculate the concentrations of strontium and hydroxyl ions and the pH of the solution.

Ans.)

Solubility of $Sr(OH)_2 = 19.23$ g/L

Then, concentration of $Sr(OH)_2$

$$= \frac{19.23}{121.63}M$$

$$= 0.1581M$$



$$\therefore [Sr^{2+}] = 0.1581M$$

$$[OH^-] = 2 \times 0.1581M = 0.3126$$

Now,

$$K_w = [OH^-][H^+] \quad \frac{10^{-14}}{0.3126} = [H^+]$$

$$\Rightarrow [H^+] = 3.2 \times 10^{-14}$$

$$\therefore pH = 13.495; 13.50$$

Q.59. The ionization constant of propanoic acid is 1.32×10^{-5} . Calculate the degree of ionization of the acid in its 0.05M solution and also its pH. What will be its degree of ionization if the solution is 0.01M in HCl also?

Ans.)

Let the degree of ionization of propanoic acid be α .

Then, representing propanoic acid as HA, we have:



$$K_\alpha = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

$$= \frac{(0.05\alpha)(0.05\alpha)}{0.05} = 0.05\alpha^2 \quad \alpha = \sqrt{\frac{K_\alpha}{0.05}} = 1.63 \times 10^{-2}$$

$$\text{Then, } [\text{H}_3\text{O}^+] = 0.05\alpha = 0.05 \times 1.63 \times 10^{-2} = K_b \cdot 15 \times 10^{-4} M$$

$$\therefore \text{pH} = 3.09$$

In the presence of 0.1M of HCl, let α' be the degree of ionization.

$$\text{Then, } [\text{H}_3\text{O}^+] = 0.01$$

$$[\text{A}^-] = 0.05 \alpha'$$

$$[\text{HA}] = 0.05$$

$$K_\alpha = \frac{0.01 \times 0.05\alpha'}{0.05}$$

$$1.32 \times 10^{-5} = 0.01 \times \alpha'$$

$$\alpha' = 1.32 \times 10^{-3}$$

Q.60. The pH of 0.1M solution of cyanic acid (HCNO) is 2.34. Calculate the ionization constant of the acid and its degree of ionization in the solution.

Ans.)

$$c = 0.1 \text{ M}$$

$$\text{pH} = 2.34$$

$$-\log [\text{H}^+] = \text{pH}$$

$$-\log [\text{H}^+] = 2.34$$

$$[\text{H}^+] = 4.5 \times 10^{-3}$$

Also,

$$[\text{H}^+] = c\alpha$$

$$4.5 \times 10^{-3} = 0.1 \times \alpha$$

$$\frac{4.5 \times 10^{-3}}{0.1} = \alpha$$

$$\alpha = 4.5 \times 10^{-3} = 0.045$$

Then,

$$K_a = c\alpha^2$$

$$= 0.1 \times (4.5 \times 10^{-3})^2$$

$$= 202.5 \times 10^{-6}$$

$$= 2.02 \times 10^{-4}$$

Q.61. The ionization constant of nitrous acid is 4.5×10^{-4} . Calculate the pH of 0.04 M sodium nitrite solution and also its degree of hydrolysis.

Ans.)

Sodium nitrite is a salt of NaOH (strong base) and HNO₂ (weak acid).

$$NO_2^- + H_2O \leftrightarrow HNO_2 + OH^- \quad K_h = \frac{[HNO_2][OH^-]}{[NO_2^-]} \Rightarrow \frac{K_w}{K_a} = \frac{10^{-14}}{4.5 \times 10^{-4}} = 22 \times 10^{-10}$$

Let, y mole of salt has undergone hydrolysis, then the concentration of various species present in the solution will be:

$$[NO_2^-] = 0.04 - y; 0.04 \quad [HNO_2] = y \quad [OH^-] = y \quad K_h = \frac{y^2}{0.04} = 0.22 \times 10^{-10} \quad y^2 =$$

$$0.0088 \times 10^{-10} \quad y = 0.093 \times 10^{-5} \therefore [OH^-] = 0.093 \times 10^{-5} M \quad [H_3O^+] =$$

$$\frac{10^{-10}}{0.093 \times 10^{-5}} = 10.75 \times 10^{-9} M$$

$$\text{Thus, } pH = -\log(10.75 \times 10^{-9})$$

$$= 7.96$$

Thus, the degree of hydrolysis is

$$= \frac{y}{0.04} = \frac{0.093 \times 10^{-5}}{0.04} = 2.325 \times 10^{-5}$$

Q.62. A 0.02M solution of pyridinium hydrochloride (C₅H₆ClN) is having pH = 3.44. Calculate the ionization constant of C₅H₅N (pyridine).

Ans.)

As we know,

$$pH = \log[H^+] \therefore [H^+] = 3.63 \times 10^{-4}$$

$$\text{Now, } K_h = \frac{3.63 \times 10^{-4}}{0.02}; \text{ (Given that concentration} = 0.02\text{M)}$$

$$\Rightarrow K_h = 6.6 \times 10^{-6}$$

As we know that,


$$K_h = \frac{K_w}{K_a} K_a = \frac{K_w}{K_h} = \frac{10^{-14}}{6.6 \times 10^{-6}}$$

$$= 1.51 \times 10^{-9}$$

Q.63 Predict if the solutions of the following salts are neutral, acidic or basic: NaCl, KBr, NaCN, NH₄NO₃, NaNO₂ and KF

Ans.)

1. KBr




Thus, it is a neutral solution.

2. NH₄NO₃



Thus, it is an acidic solution.

3. KF




Thus, it is a basic solution.

4. NaNO₂




Thus, it is a basic solution

5. NaCN



Thus, it is a basic solution.

6. NaCl



Thus, it is a neutral solution.

Q.64. The ionization constant of chloroacetic acid is 1.35×10^{-3} . What will be the pH of 0.1M acid and its 0.1M sodium salt solution?

Ans.)

The K_a for chloroacetic acid (ClCH_2COOH) is 1.35×10^{-3} .

$$\Rightarrow K_a = c\alpha^2 \therefore \alpha = \sqrt{\frac{K_a}{c}}$$

$$= \sqrt{\frac{1.35 \times 10^{-3}}{0.1}}; \text{ (given concentration = 0.1M)}$$

$$\alpha = \sqrt{1.35 \times 10^{-2}}$$

$$= 0.116$$

$$\therefore [H^+] = c\alpha = 0.1 * 0.116 = 0.0116$$

- $\text{pH} = -\log[H^+] = 1.94$

$\text{ClCH}_2\text{COONa}$ is a salt of strong base i.e. NaOH, and weak acid i.e. ClCH_2COOH



$$\text{Now, } K_h = \frac{K_w}{K_a} \quad K_h = \frac{10^{-14}}{1.35 \times 10^{-3}} = 0.740 \times 10^{-11}$$

$$\text{Also, } K_h = \frac{y^2}{0.1} \Rightarrow 0.740 \times 10^{-11} = \frac{y^2}{0.1} \Rightarrow 0.0740 \times 10^{-11} = y^2 \quad y = 0.86 \times 10^{-6}$$

$$[\text{OH}^-] = 0.86 \times 10^{-6} \therefore [H^+] = \frac{K_w}{0.86 \times 10^{-6}} = \frac{10^{-14}}{0.86 \times 10^{-6}} \quad [H^+] = 1.162 \times 10^{-3}$$

$$\text{pH} = -\log[H^+]$$

$$= 7.94$$

Q.65. Ionic product of water at 310 K is 2.7×10^{-14} . What is the pH of neutral water at this temperature?

Ans.)

Ionic Product,

$$K_w = [H^+][OH^-]$$

Assuming, $[H^+] = y$

$$\text{As, } [H^+] = [OH^-], K_w = y^2.$$

$$K_w \text{ at } 310\text{K is } 2.7 \times 10^{-14}.$$

$$\therefore 2.7 \times 10^{-14} = y^2$$

$$\bullet y = 1.64 \times 10^{-7}$$

$$\bullet [H^+] = 1.64 \times 10^{-7}$$

$$\bullet \text{pH} = -\log[H^+]$$

$$= -\log[1.64 \times 10^{-7}]$$

$$= 6.78$$

Thus, the pH of neutral water at 310K temperature is 6.78.

Q.66. Calculate the pH of the resultant mixtures:

i) 10 ml of 0.02M H_2SO_4 + 10 ml of 0.02M Ca(OH)_2

ii) 10 ml of 0.1M H_2SO_4 + 10 ml of 0.1M KOH

iii) 10 ml of 0.2M Ca(OH)_2 + 25 ml of 0.1M HCl

Ans.)

i) Moles of OH^-

$$= \frac{2 \times 10 \times 0.02}{1000} = 0.0004 \text{ mol}$$

Moles of H_3O^+

$$= \frac{2 \times 10 \times 0.02}{1000} = 0.0004 \text{ mol}$$

ii) Moles of OH^-

$$= \frac{2 \times 10 \times 0.1}{1000} = 0.002 \text{ mol}$$

Moles of H_3O^+

$$= \frac{2 \times 10 \times 0.1}{1000} = 0.001 \text{ mol}$$

Here, the H_3O^+ is in excess is 0.01 mol

$$\text{So, } [H_3O^+] = \frac{0.001}{20 \times 10^{-3}} = \frac{10^{-3}}{20 \times 10^{-3}} = 0.05$$

Thus, $\text{pH} = -\log(0.05)$

$$= 1.3$$

As the solution is neutral $\text{pH} = 7$.

iii) Moles of OH^-

$$= \frac{2 \times 10 \times 0.2}{1000} = 0.004 \text{ mol}$$

Moles of H_3O^+

$$= \frac{25 \times 0.1}{1000} = 0.0025 \text{ mol}$$

Here, the OH^- is in excess is 0.0015 mol

$$\text{So, } [OH^-] = \frac{0.0015}{35 \times 10^{-3}} = 0.0428$$

Thus, $\text{pH} = -\log(\text{OH})$

$$= 1.36$$


$$\text{pH} = 14 - 1.36 = 12.63$$

As the solution is neutral $\text{pH} = 12.63$.

Q.67. Determine the solubilities of silver chromate, barium chromate, ferric hydroxide, lead chloride and mercurous iodide at 298K from their solubility product constants given in Table 7.9. Determine also the molarities of individual ions.

Ans.)

a) Barium Chromate



$$\text{Now, } K_{\text{sp}} = [\text{Ba}^{2+}][\text{CrO}_4^{2-}]$$

Assuming the solubility of BaCrO_4 is 'x'.

Thus,

$$[\text{Ba}^{2+}] = x \text{ and } [\text{CrO}_4^{2-}] = x$$

- $K_{\text{sp}} = x^2$
- $1.2 \times 10^{-10} = x^2$
- $x = 1.09 \times 10^{-10} \text{ M}$

$$\text{Molarity of } \text{Ba}^{2+} = \text{Molarity of } \text{CrO}_4^{2-} = x = 1.09 \times 10^{-10} \text{ M}$$

b) Ferric Hydroxide



$$\text{Now, } K_{\text{sp}} = [\text{Fe}^{3+}][\text{OH}^-]^3$$

Assuming the solubility of $\text{Fe}(\text{OH})_3$ is 'x'.

Thus,

$$[\text{Fe}^{3+}] = x \text{ and } [\text{OH}^-] = 3x$$

- $K_{\text{sp}} = x(3x)^3$
- $= x \cdot 27x^3$
- $K_{\text{sp}} = 27x^4$


$$\bullet 1.0 \times 10^{-38} = 27x^4$$

$$\bullet x = 0.00037 \times 10^{-36} M$$

$$\text{Molarity of } Fe^{3+} = x = 1.39 \times 10^{-10} M$$

$$\text{Molarity of } OH^{-} = 3x = 4.17 \times 10^{-10} M$$

c) Lead Chloride



$$\text{Now, } K_{sp} = [Pb^{2+}][Cl^{-}]^2$$

Assuming the solubility of $PbCl_2$ is 'x'.

Thus,

$$[Pb^{2+}] = x \text{ and } [Cl^{-}] = 2x$$

$$\bullet K_{sp} = x(2x)^2$$

$$= x \cdot 4x^2$$

$$K_{sp} = 4x^3$$


$$\bullet 1.6 \times 10^{-5} = 4x^3$$

$$\bullet x = 1.58 \times 10^{-2} M$$

$$\text{Molarity of } Pb^{2+} = x = 1.58 \times 10^{-2} M$$

$$\text{Molarity of } Cl^{-} = 2x = 3.16 \times 10^{-2} M$$

d) Mercurous iodide



$$\text{Now, } K_{sp} = [Hg_2^{2+}][I^{-}]^2$$

Assuming the solubility of Hg_2I_2 is 'x'.

Thus,

$$[\text{Hg}^{2+}] = x \text{ and } \text{I}^{-} = 2x$$

$$\begin{aligned} \bullet K_{\text{sp}} &= x(2x)^2 \\ &= x \cdot 4x^2 \\ K_{\text{sp}} &= 4x^3 \end{aligned}$$


$$\bullet 4.5 \times 10^{-29} = 4x^3$$

$$\bullet x = 2.24 \times 10^{-10} M$$

$$\text{Molarity of } \text{Hg}^{2+} = x = 2.24 \times 10^{-10} M$$

$$\text{Molarity of } \text{I}^{-} = 2x = 4.48 \times 10^{-10} M$$

e) Silver Chromate



$$\text{Now, } K_{\text{sp}} = [\text{Ag}^{2+}]^2 [\text{CrO}_4^{2-}]$$

Assuming the solubility of Ag_2CrO_4 is 'x'.

Thus,

$$[\text{Ag}^{2+}] = 2x \text{ and } \text{CrO}_4^{2-} = x$$

$$\bullet K_{\text{sp}} = (2x)^2 \cdot x$$

$$\bullet 1.1 \times 10^{-12} = 4x^3$$


$$\bullet x = 0.65 \times 10^{-4} M$$

$$\text{Molarity of } \text{Ag}^{2+} = 2x = 1.3 \times 10^{-4} M$$

$$\text{Molarity of } \text{CrO}_4^{2-} = x = 0.65 \times 10^{-4} M$$

Q-68. The solubility product constant of Ag_2CrO_4 and AgBr are 1.1×10^{-12} and 5.0×10^{-13} respectively. Calculate the ratio of the molarities of their saturated solutions.

Ans.)



$$\text{Now, } K_{sp} = [Ag^{2+}]^2 [CrO_4^-]$$


Assuming the solubility of Ag_2CrO_4 is 'x'.

Thus,

$$[Ag^{2+}] = 2x \text{ and } [CrO_4^-] = x$$

- $K_{sp} = (2x)^2 \cdot x$
- $1.1 \times 10^{-12} = 4x^3$
- $x = 0.65 \times 10^{-4} M$

Assuming the solubility of $AgBr$ is y .



- $K_{sp} = (y)^2$
- $5.0 \times 10^{-13} = y^2$

$$y = 7.07 \times 10^{-7} M$$

The ratio of molarities to their saturated solution is:

$$\frac{x}{y} = \frac{0.65 \times 10^{-4} M}{7.07 \times 10^{-7} M} = 91.9$$

Q.69. Equal volumes of 0.002 M solutions of sodium iodate and cupric chlorate are mixed together. Will it lead to precipitation of copper iodate? (For cupric iodate $K_{sp} = 7.4 \times 10^{-8}$).

Ans.)

Cupric chlorate and sodium iodate having equal volume are mixed together, then molar concentration of cupric chlorate and sodium iodate will reduce to half.

So, molar concentration of cupric chlorate and sodium iodate in mixture is 0.001M.



$$0.0001M \qquad \qquad 0.001M$$



$$0.001M \qquad \qquad 0.001M$$

The Solubility for $Cu(IO_3)_2 \Rightarrow Cu^{2+} (aq) + 2IO_3^- (aq)$

Now, the ionic product of the copper iodate is:

$$= [\text{Cu}^{2+}] [\text{IO}_3^-]^2$$

$$= (0.001)(0.001)^2$$

$$= 1.0 \times 10^{-9} M$$

As the value of K_{sp} is more than Ionic product.

Thus, the precipitation will not occur.

Q.70. The ionization constant of benzoic acid is 6.46×10^{-5} and K_{sp} for silver benzoate is 2.5×10^{-13} . How many times is silver benzoate more soluble in a buffer of pH 3.19 compared to its solubility in pure water?

Ans.)

Here, pH = 3.19

$$[\text{H}_3\text{O}^+] = 6.46 \times 10^{-5} M$$



$$K_a \frac{[\text{C}_6\text{H}_5\text{COO}^-][\text{H}_3\text{O}^+]}{[\text{C}_6\text{H}_5\text{COOH}]} = \frac{[\text{H}_3\text{O}^+]}{K_a} = \frac{6.46 \times 10^{-4}}{6.46 \times 10^{-5}} = 10$$

Assuming the solubility of silver benzoate ($\text{C}_6\text{H}_5\text{COOAg}$) is y mol/L.

Now, $[\text{Ag}^+] = y$

$$[\text{C}_6\text{H}_5\text{COOH}] = [\text{C}_6\text{H}_5\text{COO}^-] = y$$

$$10[\text{C}_6\text{H}_5\text{COO}^-] + [\text{C}_6\text{H}_5\text{COO}^-] = y$$

$$[\text{C}_6\text{H}_5\text{COO}^-] = y/11$$

$$K_{sp}[\text{Ag}^+][\text{C}_6\text{H}_5\text{COO}^-] = y$$

$$2.5 \times 10^{-13} = y \frac{y}{11}$$

$$y = 1.66 \times 10^{-6} \text{ mol/L}$$

Hence, solubility of $\text{C}_6\text{H}_5\text{COOAg}$ in buffer of pH = 3.19 is 1.66×10^{-6} mol/L.

For, water:

Assuming the solubility of silver benzoate ($\text{C}_6\text{H}_5\text{COOAg}$) is x mol/L.

Now, $[Ag^+] = x \text{ M}$

$$K_{sp} = [Ag^+][C_6H_5COO^-]$$

$$K_{sp} = (y)^2$$

$$y = \sqrt{K_{sp}} = \sqrt{2.5 \times 10^{-13}} = 5 \times 10^{-7} \text{ mol/L} \therefore \frac{y}{x} = \frac{1.66 \times 10^{-6}}{5 \times 10^{-7}} = 3.32$$

Thus, the solubility of silver benzoate in water is 3.32 times the solubility of silver benzoate in pH = 3.19.

Q.71. What is the maximum concentration of equimolar solutions of ferrous sulphate and sodium sulphide so that when mixed in equal volumes, there is no precipitation of iron sulphide? (For iron sulphide, $K_{sp} = 6.3 \times 10^{-18}$).

Ans.)


Assuming the maximum concentration of each solution is $y \text{ mol/L}$

On mixing the solutions the volume of the concentration of each solution is reduced to half.

After mixing the maximum concentration of each solution is $y/2 \text{ mol/L}$.

$$\text{Thus, } [FeSO_4] = [Na_2S] = y/2 \text{ M}$$

$$\text{So, } [Fe^{2+}] = [FeSO_4] = y/2 \text{ M}$$




$$K_{sp} = [Fe^{2+}][S^{2-}] \quad 6.3 \times 10^{-18} = \left(\frac{y}{2}\right)\left(\frac{y}{2}\right) \frac{y^2}{4} = 6.3 \times 10^{-18}$$

$$\text{Thus, } y = 5.02 \times 10^{-9}$$

Thus, if the concentration of $FeSO_4$ and Na_2SO_4 are equal to or less than that of $5.02 \times 10^{-9} \text{ M}$, then there won't be precipitation of FeS .

Q.72. What is the minimum volume of water required to dissolve 1g of calcium sulphate at 298 K? (For calcium sulphate, K_{sp} is 9.1×10^{-6}).

Ans.)



$$K_{sp} = [Ca^{2+}][SO_4^{2-}]$$

Assuming the solubility of calcium sulphate is x .

$$\text{So, } K_{sp} = x^2$$

$$\therefore 9.1 \times 10^{-6} = x^2 \therefore x = 3.02 \times 10^{-3} \text{ mol/L}$$

Now, molecular mass of calcium sulphate is 136g/mol.

Solubility in calcium sulphate in g/mol is

$$= 3.02 \times 10^{-3} \times 136$$

$$= 0.41 \text{ g/L}$$

i.e. 1 litre H₂O will be required to dissolve 0.41g of calcium sulphate.

Thus, minimum volume of H₂O required to dissolve 1 gram of CaSO₄ at 298K is

$$= \frac{1}{0.41} L = 2.44L$$

Q.73. The concentration of sulphide ion in 0.1M HCl solution saturated with hydrogen sulphide is $1.0 \times 10^{-19} \text{ M}$. If 10 mL of this is added to 5 mL of 0.04 M solution of the following: FeSO₄, MnCl₂, ZnCl₂ and CdCl₂. In which of these solutions precipitation will take place?

Ans.)

If the ionic product exceeds the K_{sp} value, then only precipitation can take place.

Before mixing:

$$[S^{2-}] = K_{sp} = 1.0 \times 10^{-19} \text{ M} \quad [M^{2+}] = 0.04 \text{ M}$$

Volume = 10mL

Volume = 5mL

After mixing:

$$[S^{2-}] = ? \text{ and}$$

$$[M^{2+}] = ?$$

Total volume = (10 + 5) = 15mL

Volume = 15mL

$$[S^{2-}] = \frac{1.0 \times 10^{-19} \times 10}{15} = 6.67 \times 10^{-20} \text{ M} \quad [M^{2+}] = \frac{0.04 \times 5}{15} = 1.33 \times 10^{-2} \text{ M}$$

Now, the ionic product = $[M^{2+}][S^{2-}]$

$$= (1.33 \times 10^{-2})(6.67 \times 10^{-20})$$

$$= 8.87 \times 10^{-22}$$

Here, the ionic product of CdS and ZnS exceeds its corresponding K_{sp} value.

Thus, precipitation will occur in ZnCl₂ and CdCl₂ solutions.